

FORCED WAVES FOR A THREE-SPECIES PREDATOR-PREY SYSTEM WITH NONLOCAL DISPERSAL IN A SHIFTING ENVIRONMENT

JONG-SHENQ GUO, FRANÇOIS HAMEL, AND CHIN-CHIN WU

ABSTRACT. We consider a three-species predator-prey system involving two competing predators and one prey. The species diffuse with nonlocal dispersal kernels with possibly non-compact support, and they interact in a heterogeneous environment moving with a positive forced speed such that the environment is favorable to the prey in the absence of predators far ahead of the shifting boundary and it is unfavorable far behind. Such systems arise in the modeling of population dynamics under the effect of a shifting environment, such as climate change. We show on the one hand the existence of waves connecting the trivial state to the unique constant positive co-existence state for any value of the forced speed. On the other hand, we show the existence of critical positive speeds for the existence of waves connecting the trivial state to the states corresponding to the absence of one or two predators.

1. INTRODUCTION

Nonlocal dispersal models arise in population dynamics to describe long-distance dispersal of individuals [24, 25, 28]. Recently, the effect of environmental heterogeneity has drawn a lot of attention in biological applications. In this paper, we consider the following predator-prey system with two weak competing predators and one prey:

$$(1.1) \quad \begin{cases} u_t(x, t) &= d_1 \mathcal{N}_1[u](x, t) + r_1 u(x, t) [-1 - u(x, t) - kv(x, t) + aw(x, t)], \\ v_t(x, t) &= d_2 \mathcal{N}_2[v](x, t) + r_2 v(x, t) [-1 - hu(x, t) - v(x, t) + aw(x, t)], \\ w_t(x, t) &= d_3 \mathcal{N}_3[w](x, t) + r_3 w(x, t) [\alpha(x - st) - bu(x, t) - bv(x, t) - w(x, t)], \end{cases}$$

in which the nonnegative quantities $u = u(x, t)$ and $v = v(x, t)$ are the densities of two predators and $w = w(x, t)$ is the density of the single prey, for $x, t \in \mathbb{R}$.

Throughout the paper, the parameters $d_1, d_2, d_3, r_1, r_2, r_3, a, b, h, k$ are all positive, and a, b, h, k are such that

$$(1.2) \quad a > 1, \quad 0 < h, k < 1, \quad 0 < b < \frac{1}{2(a-1)}.$$

Date: November 17, 2022. Corresponding author: C.-C. Wu.

This work was supported in part by the Ministry of Science and Technology of Taiwan under the grants 108-2115-M-032-006-MY3 (JSG) and 110-2115-M-005-001-MY2 (CCW), and by the CNRS-NCTS joint International Research Network ReaDiNet. This work has also received funding from Excellence Initiative of Aix-Marseille Université - A*MIDEX, a French “Investissements d’Avenir” programme, and from the ANR RESISTE (ANR-18-CE45-0019) project.

Key words and phrases. Predator-prey model, nonlocal dispersal, climate change, forced wave.

Meanwhile, $\mathcal{N}_1[u]$, $\mathcal{N}_2[v]$ and $\mathcal{N}_3[w]$ formulate the spatial nonlocal dispersal and are defined by

$$\begin{cases} \mathcal{N}_1[u](x, t) & := (J_1 * u)(x, t) - u(x, t) = \int_{\mathbb{R}} J_1(y)u(x-y, t)dy - u(x, t), \\ \mathcal{N}_2[v](x, t) & := (J_2 * v)(x, t) - v(x, t) = \int_{\mathbb{R}} J_2(y)v(x-y, t)dy - v(x, t), \\ \mathcal{N}_3[w](x, t) & := (J_3 * w)(x, t) - w(x, t) = \int_{\mathbb{R}} J_3(y)w(x-y, t)dy - w(x, t), \end{cases}$$

in which, throughout the paper, the functions $J_i : \mathbb{R} \rightarrow \mathbb{R}$ ($i = 1, 2, 3$) are probability kernel functions satisfying

- (J1) J_i is nonnegative, measurable with respect to the Lebesgue measure, $\int_{\mathbb{R}} J_i(y)dy = 1$, and there are $\eta_i > 0$ and $y_i^\pm \in \mathbb{R}$ such that $y_i^- < 0 < y_i^+$ and $J_i > 0$ in $(y_i^- - \eta_i, y_i^- + \eta_i)$ and in $(y_i^+ - \eta_i, y_i^+ + \eta_i)$;
- (J2) $\int_{\mathbb{R}} J_i(y)y dy = 0$;
- (J3) there exist $-\infty \leq \tilde{\lambda}_i < 0 < \hat{\lambda}_i \leq +\infty$ such that $I_i(\lambda) < +\infty$ for all $\lambda \in (\tilde{\lambda}_i, \hat{\lambda}_i)$, and $I_i(\lambda) \rightarrow +\infty$ as $\lambda \downarrow \tilde{\lambda}_i$ and $\lambda \uparrow \hat{\lambda}_i$, where

$$I_i(\lambda) := \int_{\mathbb{R}} J_i(y)e^{\lambda y} dy.$$

Conditions (J1)-(J3) imply that each function I_i is strictly convex and of class C^∞ in $(\tilde{\lambda}_i, \hat{\lambda}_i)$, that $I_i(0) = 1$, $I_i'(0) = 0$, and $I_i(\lambda) = +\infty$ for all $\lambda \in (-\infty, \tilde{\lambda}_i]$ if $-\infty < \tilde{\lambda}_i$ (resp. for all $\lambda \in [\hat{\lambda}_i, +\infty)$ if $\hat{\lambda}_i < +\infty$). If J_i has a compact support, then $\tilde{\lambda}_i = -\infty$ and $\hat{\lambda}_i = +\infty$. Conditions (J1) and (J3) imply that each function $y \mapsto J_i(y)y$ is in $L^1(\mathbb{R})$ and, if J_i is even, they necessarily yield (J2), and $\tilde{\lambda}_i = -\hat{\lambda}_i$ in (J3). We point out that the conditions (J1)-(J3) are satisfied in particular if J_i is nonnegative, continuous, even, has a unit integral over \mathbb{R} and if $I_i(\lambda) < +\infty$ for some $\lambda > 0$. But the conditions (J1)-(J3) cover more general dispersal kernels J_i , in particular the kernels J_i can be non-symmetric.

The function α in (1.1), describing the heterogeneity, is assumed to be continuous in \mathbb{R} and the given positive constant s denotes the environmental shifting speed. Throughout this paper we also impose the following conditions on α :

- ($\alpha 1$) α has limits $\alpha(\pm\infty)$ at $\pm\infty$, such that $-\infty < \alpha(-\infty) < 0 < \alpha(+\infty) < +\infty$, and $\alpha(z) \leq \alpha(+\infty)$ for all $z \in \mathbb{R}$;
- ($\alpha 2$) there exist $C > 0$ and $\rho > 0$ such that $\alpha(+\infty) - \alpha(z) \leq Ce^{-\rho z}$ for all large z .

Condition ($\alpha 1$) means that the environment is favourable to the prey ahead of the shifting boundary $x = st$, then gradually deteriorates until it becomes hostile to the species far behind this shifting boundary. We point out that the function α is not assumed to be monotone.

Biologically, parameters (d_1, d_2, d_3) , (h, k) , a and b represent the diffusion coefficients, competition rates, conversion rate and predation rate, respectively. The assumption $0 < h, k < 1$ means that the two predators are weak competitors. Moreover, the negative net growth rate $-r_i$ ($i = 1, 2$) of each predator means that each predator cannot survive without the feeding of the prey. The terms $-r_1 u(x, t)^2$ and $-r_2 v(x, t)^2$ stand for the intra-specific competition inside each predator population. The smallness of the predation rate b means

that the predation has a relatively low impact on the prey, whereas the largeness of the conversion rate a is related to the relatively large fitness of the predators in the presence of the prey. The intrinsic growth rate of prey is given by $r_3\alpha(x - st)$ which is temporal-spatial dependent and takes both positive and negative values. From the modeling point of view, the function α represents the shifting environment effect, such as climate change. Although this term does not appear in the equation of each predator, the changing effect actually affects indirectly both predators due to the fact that predators are fed by prey.

We are concerned with the existence of forced waves for (1.1). Namely, a traveling wave solution of (1.1) with speed s is a solution in the form

$$(u, v, w)(x, t) = (\phi_1, \phi_2, \phi_3)(z), \quad z := x - st,$$

with $C^1(\mathbb{R})$ functions ϕ_1, ϕ_2, ϕ_3 . Then (ϕ_1, ϕ_2, ϕ_3) satisfies

$$(1.3) \quad \begin{cases} -s\phi_1'(z) = d_1\mathcal{N}_1[\phi_1](z) + r_1\phi_1(z) [-1 - \phi_1(z) - k\phi_2(z) + a\phi_3(z)], & z \in \mathbb{R}, \\ -s\phi_2'(z) = d_2\mathcal{N}_2[\phi_2](z) + r_2\phi_2(z) [-1 - h\phi_1(z) - \phi_2(z) + a\phi_3(z)], & z \in \mathbb{R}, \\ -s\phi_3'(z) = d_3\mathcal{N}_3[\phi_3](z) + r_3\phi_3(z) [\alpha(z) - b\phi_1(z) - b\phi_2(z) - \phi_3(z)], & z \in \mathbb{R}, \end{cases}$$

where

$$\mathcal{N}_i[\phi_i](z) := \int_{\mathbb{R}} J_i(y)\phi_i(z - y)dy - \phi_i(z), \quad i = 1, 2, 3.$$

Throughout the paper, by a solution of (1.3), we always mean a triplet (ϕ_1, ϕ_2, ϕ_3) of $C^1(\mathbb{R})$ nonnegative bounded functions. Since the environment is hostile to the species far behind the shifting boundary $x = st$, by the assumption on α , it can be expected that all species go extinction eventually. Hence we impose the boundary condition

$$(\phi_1, \phi_2, \phi_3)(-\infty) = (0, 0, 0).$$

On the other hand, without loss of generality we may assume that

$$(1.4) \quad \alpha(+\infty) = 1.$$

Condition (1.4) is assumed in all main results (Theorems 1.1-1.6). Then the following states are the only possible constant and non-trivial limiting states $(\phi_1, \phi_2, \phi_3)(+\infty)$ at $z = +\infty$:

$$E_1 := (0, 0, 1), \quad E_2 := (u_p, 0, w_p), \quad E_3 := (0, u_p, w_p), \quad E_4 := (u^*, v^*, w^*),$$

where the real numbers u_p, w_p, u^*, v^*, w^* are all positive and given by

$$\begin{cases} u_p := \frac{a-1}{ab+1}, \quad w_p := \frac{b+1}{ab+1}, \\ w^* := \frac{1+b\gamma}{1+ab\gamma}, \quad \gamma := \frac{2-h-k}{1-hk} > 1, \quad v^* := \frac{1-h}{1-hk}(aw^* - 1), \quad u^* := \frac{1-k}{1-hk}(aw^* - 1). \end{cases}$$

Biologically, the state E_1 corresponds to a saturated aboriginal prey living in the habitat and there are two invading alien predators; E_2 or E_3 is a pair of aboriginal co-existent predator-prey and an invading alien predator; lastly, E_4 is the positive co-existence state.

The study of forced waves has attracted a lot of attention recently. We refer the reader to, e.g., [2, 3, 4, 5, 6, 18, 19, 22, 23, 29, 38] for the case of scalar equations with local diffusion and to [12, 13, 27, 31, 37] for the case of scalar equations with nonlocal dispersal.

For two-species models, we refer the reader to [34] for a cooperative model, [1, 14, 30, 33, 35] for competition models, and [10] for a predator-prey system with both classical diffusion and nonlocal dispersal. Recently, forced waves for a three-species predator-prey system were investigated in [11] for two competing preys and one predator with the classical diffusion. In [11], the authors obtained both front and pulse types forced waves.

In the case of nonlocal dispersal, one usually assume that each kernel function is of compact support (so that $\tilde{\lambda}_i = -\infty$ and $\hat{\lambda}_i = +\infty$ in (J3)). One of the motivations of this work is to remove the restriction on the compact support of kernel(s). This question has been left open in [10] for the two-species predator-prey system.

We now describe our main results as follows. We repeat that the condition (1.2) is assumed in all results, as are (J1)-(J3) and ($\alpha 1$)-($\alpha 2$). First, the following theorem provides the existence of waves connecting $(0, 0, 0)$ and the co-existence state E_4 , whatever the positive forced speed s may be.

Theorem 1.1. *In addition to (1.2) and (1.4), suppose*

$$(1.5) \quad b < \min \left\{ \frac{1-h}{2a}, \frac{1-k}{2a} \right\}.$$

Then, for any $s > 0$, there exists a positive¹ solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) such that

$$(\phi_1, \phi_2, \phi_3)(-\infty) = (0, 0, 0) \quad \text{and} \quad (\phi_1, \phi_2, \phi_3)(+\infty) = E_4.$$

Next, for waves connecting $(0, 0, 0)$ and the predator-free state $E_1 = (0, 0, 1)$, we let

$$(1.6) \quad s_i^* := \inf_{\lambda \in (0, \hat{\lambda}_i)} Q_i(\lambda), \quad Q_i(\lambda) := \frac{d_i[I_i(\lambda) - 1] + r_i(a-1)}{\lambda}, \quad \lambda \in (0, \hat{\lambda}_i), \quad i = 1, 2.$$

From (1.2) and (J1)-(J3) and the comments after (J1)-(J3), each function Q_i ($i = 1, 2$) is continuous and positive in $(0, \hat{\lambda}_i)$, $Q_i(\lambda) \rightarrow +\infty$ as $\lambda \downarrow 0$ or $\lambda \uparrow \hat{\lambda}_i$ (this last property is immediate if $\hat{\lambda}_i < +\infty$, and it holds as well if $\hat{\lambda}_i = +\infty$ since $I_i'(\lambda) = \int_{\mathbb{R}} J_i(y) y e^{\lambda y} dy \rightarrow +\infty$ as $\lambda \rightarrow +\infty = \hat{\lambda}_i$ in that case). Hence, each s_i^* is positive and the infimum in (1.6) is a minimum, and it is reached by a unique $\lambda_i^* \in (0, \hat{\lambda}_i)$ (since I_i'' is positive in $(0, \hat{\lambda}_i)$, and even in the whole interval of definition $(\tilde{\lambda}_i, \hat{\lambda}_i)$). Furthermore, Q_i is decreasing in $(0, \lambda_i^*]$ and increasing in $[\lambda_i^*, \hat{\lambda}_i)$. Then we have

Theorem 1.2. *Suppose (1.2) and (1.4). If $s > \max\{s_1^*, s_2^*\}$, then there exists a positive solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) such that*

$$(1.7) \quad (\phi_1, \phi_2, \phi_3)(-\infty) = (0, 0, 0) \quad \text{and} \quad (\phi_1, \phi_2, \phi_3)(+\infty) = E_1.$$

Moreover, when $(\tilde{\lambda}_i, \hat{\lambda}_i) = (-\infty, +\infty)$ for some $i \in \{1, 2\}$, then a positive solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) and (1.7) exists only if $s \geq s_i^$.*

For the waves connecting $(0, 0, 0)$ and the mixed state $E_2 = (u_p, 0, w_p)$, we let

$$(1.8) \quad \beta_2 := -1 - hu_p + aw_p = \frac{(a-1)(1-h)}{ab+1} > 0$$

¹By positive, we mean that each component ϕ_i ($i = 1, 2, 3$) is positive in \mathbb{R} .

and

$$(1.9) \quad s_2^{**} := \inf_{\lambda \in (0, \hat{\lambda}_2)} R_2(\lambda), \quad R_2(\lambda) := \frac{d_2[I_2(\lambda) - 1] + r_2\beta_2}{\lambda}, \quad \lambda \in (0, \hat{\lambda}_2).$$

As for Q_i in (1.6), the function R_2 is continuous and positive in $(0, \hat{\lambda}_2)$, $R_2(\lambda) \rightarrow +\infty$ as $\lambda \downarrow 0$ or $\lambda \uparrow \hat{\lambda}_2$, s_2^{**} is positive and the infimum in the definition of s_2^{**} is a minimum, reached by a unique $\lambda_2^{**} \in (0, \hat{\lambda}_2)$. Furthermore, R_2 is decreasing in $(0, \lambda_2^{**}]$ and increasing in $[\lambda_2^{**}, \hat{\lambda}_2)$. Then we have

Theorem 1.3. *In addition to (1.2) and (1.4), suppose*

$$(1.10) \quad \max\{d_1, d_3\} \leq d_2, \quad J_1 = J_2 = J_3 \text{ in } \mathbb{R},$$

and

$$(1.11) \quad r_1[1 + k(a - 1)] \leq r_2\beta_2.$$

If $s > s_2^{**}$ and $s \geq R_2(\rho)$, with $\rho > 0$ as in $(\alpha 2)$, then there exists a positive solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) such that

$$(1.12) \quad (\phi_1, \phi_2, \phi_3)(-\infty) = (0, 0, 0) \quad \text{and} \quad (\phi_1, \phi_2, \phi_3)(+\infty) = E_2.$$

Moreover, when $(\tilde{\lambda}_2, \hat{\lambda}_2) = (-\infty, +\infty)$, a positive solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) and (1.12) exists only if $s \geq s_2^{**}$.

Notice that, if condition $(\alpha 2)$ is satisfied for a certain $\rho > 0$, so is it for every $\rho' \in (0, \rho]$. Therefore, in Theorem 1.3, one can always assume without loss of generality that $\rho \in (0, \hat{\lambda}_2)$ and that $R_2(\rho)$ is a real number. We also point out that, if $\alpha(z) = \alpha(+\infty)$ for all z large enough, then $(\alpha 2)$ is satisfied for all $\rho > 0$, hence ρ can be chosen so that $R_2(\rho) = s_2^{**}$ and the condition $s \geq R_2(\rho)$ in the statement can then be dropped.

By exchanging the roles of u and v , a similar theorem to Theorem 1.3, on forced waves connecting the trivial state and E_3 , can be derived. We omit it here.

One should notice that the above theorems hold under the assumptions (J1)-(J3) on kernels J_i , $i = 1, 2, 3$, and conditions $(\alpha 1)$ - $(\alpha 2)$ on α . In particular, the kernel functions J_i , $i = 1, 2, 3$, are not assumed to be compactly supported or symmetric. It is worth to mention that the method of this work can also be applied to the 2-species system studied in [10] and that the compact support assumption and the symmetry of the kernels can then be removed.

Next, we turn to the derivation of waves with critical speeds. We divide our main results into three cases. The following two theorems show, under some conditions, the existence of forced waves connecting the trivial state to the predator-free state $E_1 = (0, 0, 1)$, for the minimal possible shifting speed. We recall that s_1^* and s_2^* are defined in (1.6).

Theorem 1.4. *In addition to (1.2) and (1.4), suppose that $s_1^* = s_2^*$, and that J_1 and J_2 have compact supports. Then a positive solution of (1.3) with (1.7) exists for $s = s_1^* = s_2^*$.*

Note that the condition $s_1^* = s_2^*$ is fulfilled especially if $d_1 = d_2$, $r_1 = r_2$, and $J_1 = J_2$ in \mathbb{R} .

Theorem 1.5. *In addition to (1.2) and (1.4), suppose that $s_1^* > s_2^*$ and that J_1 has a compact support. Then a positive solution of (1.3) with (1.7) exists for $s = s_1^*$.*

Note that, from the comments after (J1)-(J3), the condition $s_1^* > s_2^*$ is fulfilled if $d_1 \geq d_2$, $r_1 \geq r_2$, $(d_1, r_1) \neq (d_2, r_2)$, and $J_1 = J_2$ in \mathbb{R} . A similar result to Theorem 1.5 also holds if $s_2^* > s_1^*$ and if J_2 has a compact support (namely, in that case, a forced wave connecting the trivial state and E_1 with the critical speed $s = s_2^*$ exists).

Lastly, the next theorem shows, under some conditions, the existence of forced waves connecting the trivial state to the mixed state $E_2 = (u_p, 0, w_p)$, for the minimal possible shifting speed.

Theorem 1.6. *In addition to (1.2) and (1.4), suppose that $d_1 = d_2 = d_3$, that $J := J_1 = J_2 = J_3$ in \mathbb{R} , and that J has a compact support. We also assume that (1.11) holds and that $\rho \geq \lambda_2^{**}$, where $\rho > 0$ is as in ($\alpha 2$) and $\lambda_2^{**} > 0$ denotes the unique minimum of R_2 in $(0, \hat{\lambda}_2) = (0, +\infty)$. Then a positive solution of (1.3) with (1.12) exists for $s = s_2^{**}$.*

By exchanging the roles of u and v , a similar theorem to Theorem 1.6 on forced waves connecting the trivial state and E_3 with critical speed can be derived. We omit it here.

Predator-prey systems such as (1.1) are neither cooperative nor competitive, and the maximum principle does not hold for these systems. Furthermore, due to the heterogeneity of the function α , the systems (1.1) and (1.3) are not invariant by translation with respect to the spatial variable, and new difficulties then arise in comparison with the existence of traveling waves for homogeneous predator-prey systems (we refer to, e.g., [7, 8, 9, 20, 21] for the existence of waves for various homogeneous predator-prey systems with local diffusion, and to [15, 16, 17] for the study of spreading speeds for such homogeneous systems with local or nonlocal dispersal).

With respect to the paper [10] on systems with both classical diffusion and nonlocal dispersal, we introduce in the present paper, in order to remove the assumption on compact support, some new ideas based on the Schauder fixed-point theorem, the suitable definition of upper and lower solutions, and translation strategies described in Section 2. The existence of forced waves connecting co-existence or predator-free states to the trivial state then relies on the construction of suitable upper and lower solutions, which is quite intricate for this three-species system.

The question of existence of traveling waves or forced waves has been addressed extensively. For the stability of traveling waves, or the convergence of general solutions to traveling waves, we refer the reader to [31] and the references cited therein for scalar equations or systems which admit a comparison principle. However, without comparison principle, the question of convergence to traveling waves is still open.

Organization of the paper. Section 2 presents a general framework for the existence of waves, based on fixed point methods and translation strategies. Theorems 1.1-1.3 on the existence of super-critical waves connecting $(0, 0, 0)$ and E_4 , E_1 or E_2 are proved in Sections 3-5. Section 6 is devoted to the proof of Theorems 1.4-1.6 on the existence of forced waves with critical speeds.

2. PRELIMINARIES

First, we introduce the notion of generalized upper-lower solutions of (1.3).

Definition 2.1. Continuous functions $(\bar{\phi}_1, \bar{\phi}_2, \bar{\phi}_3)$ and $(\underline{\phi}_1, \underline{\phi}_2, \underline{\phi}_3)$ are called a pair of upper and lower solutions of (1.3) if $\bar{\phi}_i \geq \underline{\phi}_i$ in \mathbb{R} , $i = 1, 2, 3$, and the following inequalities

$$(2.1) \quad \mathcal{U}_1(z) := d_1 \mathcal{N}_1[\bar{\phi}_1](z) + s\bar{\phi}_1'(z) + r_1 \bar{\phi}_1(z) [-1 - \bar{\phi}_1(z) - k\underline{\phi}_2(z) + a\bar{\phi}_3(z)] \leq 0,$$

$$(2.2) \quad \mathcal{U}_2(z) := d_2 \mathcal{N}_2[\bar{\phi}_2](z) + s\bar{\phi}_2'(z) + r_2 \bar{\phi}_2(z) [-1 - h\underline{\phi}_1(z) - \bar{\phi}_2(z) + a\bar{\phi}_3(z)] \leq 0,$$

$$(2.3) \quad \mathcal{U}_3(z) := d_3 \mathcal{N}_3[\bar{\phi}_3](z) + s\bar{\phi}_3'(z) + r_3 \bar{\phi}_3(z) [\alpha(z) - b\underline{\phi}_1(z) - b\underline{\phi}_2(z) - \bar{\phi}_3(z)] \leq 0,$$

$$(2.4) \quad \mathcal{L}_1(z) := d_1 \mathcal{N}_1[\underline{\phi}_1](z) + s\underline{\phi}_1'(z) + r_1 \underline{\phi}_1(z) [-1 - \underline{\phi}_1(z) - k\bar{\phi}_2(z) + a\underline{\phi}_3(z)] \geq 0,$$

$$(2.5) \quad \mathcal{L}_2(z) := d_2 \mathcal{N}_2[\underline{\phi}_2](z) + s\underline{\phi}_2'(z) + r_2 \underline{\phi}_2(z) [-1 - h\bar{\phi}_1(z) - \underline{\phi}_2(z) + a\underline{\phi}_3(z)] \geq 0,$$

$$(2.6) \quad \mathcal{L}_3(z) := d_3 \mathcal{N}_3[\underline{\phi}_3](z) + s\underline{\phi}_3'(z) + r_3 \underline{\phi}_3(z) [\alpha(z) - b\bar{\phi}_1(z) - b\bar{\phi}_2(z) - \underline{\phi}_3(z)] \geq 0$$

hold for all $z \in \mathbb{R} \setminus E$ for some finite subset E (possibly empty) of \mathbb{R} (the functions $\underline{\phi}_i$ and $\bar{\phi}_i$ are assumed to be of class C^1 in, at least, $\mathbb{R} \setminus E$).

Next, from the Schauder fixed-point theorem, we can derive the existence of wave profiles as follows. Note that our proof below actually only requires the nonnegativity and integrability of the kernels J_i .

Lemma 2.2. Let $s > 0$ be given. Let $(\bar{\phi}_1, \bar{\phi}_2, \bar{\phi}_3)$ and $(\underline{\phi}_1, \underline{\phi}_2, \underline{\phi}_3)$ be a pair of nonnegative bounded upper and lower solutions of (1.3). Then (1.3) admits a solution (ϕ_1, ϕ_2, ϕ_3) such that $\underline{\phi}_i \leq \phi_i \leq \bar{\phi}_i$ in \mathbb{R} for all $i = 1, 2, 3$.

Proof. First of all, we denote

$$M := \max \{ \|\alpha\|_{L^\infty(\mathbb{R})}, \|\bar{\phi}_1\|_{L^\infty(\mathbb{R})}, \|\bar{\phi}_2\|_{L^\infty(\mathbb{R})}, \|\bar{\phi}_3\|_{L^\infty(\mathbb{R})} \}$$

and we define a positive constant β as follows:

$$(2.7) \quad \beta := \max\{\sigma_1, \sigma_2, \sigma_3\} > 0,$$

where

$$(2.8) \quad \sigma_1 := d_1 + r_1(2M + kM + 1), \quad \sigma_2 := d_2 + r_2(2M + hM + 1), \quad \sigma_3 := d_3 + r_3M(b + 3).$$

We also fix an arbitrary positive continuous function $W : \mathbb{R} \rightarrow \mathbb{R}$ (a weight function) such that $W(z) \rightarrow 0$ as $z \rightarrow \pm\infty$ and we consider the Banach space

$$X := \{(\phi_1, \phi_2, \phi_3) : \phi_i \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R}), i = 1, 2, 3\}$$

endowed with the norm

$$(2.9) \quad \|(\phi_1, \phi_2, \phi_3)\|_X := \max \{ \|\phi_1 W\|_{L^\infty(\mathbb{R})}, \|\phi_2 W\|_{L^\infty(\mathbb{R})}, \|\phi_3 W\|_{L^\infty(\mathbb{R})} \}.$$

We then set

$$\Gamma := \{(\phi_1, \phi_2, \phi_3) : \phi_i \in C^0(\mathbb{R}), 0 \leq \underline{\phi}_i \leq \phi_i \leq \bar{\phi}_i \text{ in } \mathbb{R}, i = 1, 2, 3\},$$

which is a non-empty convex closed bounded subset of $(X, \|\cdot\|_X)$.

Then we transform the differential system (1.3) to an integral operator (as in e.g. [17, 21]), and we will apply a fixed-point theorem in Γ . To do so, we set, for $\Phi := (\phi_1, \phi_2, \phi_3) \in \Gamma$,

$$\begin{cases} F_1[\Phi](y) := \beta\phi_1(y) + d_1\mathcal{N}_1[\phi_1](y) + r_1\phi_1(y) [-1 - \phi_1(y) - k\phi_2(y) + a\phi_3(y)], & y \in \mathbb{R}, \\ F_2[\Phi](y) := \beta\phi_2(y) + d_2\mathcal{N}_2[\phi_2](y) + r_2\phi_2(y) [-1 - h\phi_1(y) - \phi_2(y) + a\phi_3(y)], & y \in \mathbb{R}, \\ F_3[\Phi](y) := \beta\phi_3(y) + d_3\mathcal{N}_3[\phi_3](y) + r_3\phi_3(y) [\alpha(y) - b\phi_1(y) - b\phi_2(y) - \phi_3(y)], & y \in \mathbb{R}, \end{cases}$$

and then

$$P[\Phi] := (P_1[\Phi], P_2[\Phi], P_3[\Phi]),$$

where

$$P_i[\Phi](z) := \frac{e^{\beta z/s}}{s} \int_z^{+\infty} e^{-\beta y/s} F_i[\Phi](y) dy = \frac{1}{s} \int_0^{+\infty} e^{-\beta u/s} F_i[\Phi](z+u) du, \quad z \in \mathbb{R}, \quad i = 1, 2, 3.$$

For each $\Phi \in \Gamma$, the functions $F_i[\Phi]$ are bounded and continuous in \mathbb{R} and then so are the functions $P_i[\Phi]$, for $i = 1, 2, 3$. Furthermore, the functions $P_i[\Phi]$ are actually of class $C^1(\mathbb{R})$. It is straightforward to check that a fixed point $\Phi = (\phi_1, \phi_2, \phi_3)$ of P in Γ is a $C^1(\mathbb{R})$ solution of (1.3) such that $\underline{\phi}_i \leq \phi_i \leq \bar{\phi}_i$ in \mathbb{R} for $i = 1, 2, 3$.

To derive a fixed point of P in Γ , we first show that P maps Γ into Γ . To this aim, we let $\Phi \in \Gamma$. Then, using $\beta \geq \sigma_1$ in (2.7)-(2.8) together with the definition of Γ , we get that

$$F_1[\Phi](z) \geq F_1[(\underline{\phi}_1, \bar{\phi}_2, \underline{\phi}_3)](z),$$

hence $P_1[\Phi](z) \geq P_1[(\underline{\phi}_1, \bar{\phi}_2, \underline{\phi}_3)](z)$, for all $z \in \mathbb{R}$. On the other hand, by the definition of upper-lower solutions, especially (2.4) and the finiteness of the set E in Definition 2.1, we have

$$\begin{aligned} P_1[(\underline{\phi}_1, \bar{\phi}_2, \underline{\phi}_3)](z) &= \frac{1}{s} \int_z^{+\infty} e^{\beta(z-y)/s} F_1[(\underline{\phi}_1, \bar{\phi}_2, \underline{\phi}_3)](y) dy \\ &= \lim_{\varepsilon \downarrow 0} \int_{[z, +\infty) \setminus \cup_{t \in E} (t-\varepsilon, t+\varepsilon)} e^{\beta(z-y)/s} \frac{F_1[(\underline{\phi}_1, \bar{\phi}_2, \underline{\phi}_3)](y)}{s} dy \\ &\geq \liminf_{\varepsilon \downarrow 0} \int_{[z, +\infty) \setminus \cup_{t \in E} (t-\varepsilon, t+\varepsilon)} e^{\beta(z-y)/s} \left(-\underline{\phi}'_1(y) + \frac{\beta}{s} \underline{\phi}_1(y) \right) dy = \underline{\phi}_1(z) \end{aligned}$$

for all $z \in \mathbb{R}$, since $\underline{\phi}_1$ is continuous in \mathbb{R} . Hence, $P_1[\Phi](z) \geq \underline{\phi}_1(z)$ for all $z \in \mathbb{R}$. Similarly, we can derive

$$\begin{cases} P_1[\Phi] \leq P_1[(\bar{\phi}_1, \underline{\phi}_2, \bar{\phi}_3)] \leq \bar{\phi}_1 & \text{in } \mathbb{R}, \\ \underline{\phi}_2 \leq P_2[(\bar{\phi}_1, \underline{\phi}_2, \underline{\phi}_3)] \leq P_2[\Phi] \leq P_2[(\underline{\phi}_1, \bar{\phi}_2, \bar{\phi}_3)] \leq \bar{\phi}_2 & \text{in } \mathbb{R}, \\ \underline{\phi}_3 \leq P_3[(\bar{\phi}_1, \bar{\phi}_2, \underline{\phi}_3)] \leq P_3[\Phi] \leq P_3[(\underline{\phi}_1, \underline{\phi}_2, \bar{\phi}_3)] \leq \bar{\phi}_3 & \text{in } \mathbb{R}, \end{cases}$$

by the choice of β in (2.7)-(2.8) and the definition of upper-lower solutions. Hence, we finally obtain that $P(\Gamma) \subset \Gamma$.

Now, we observe that, from the definition of Γ and the boundedness of the functions $\underline{\phi}_i$ and $\bar{\phi}_i$ ($i = 1, 2, 3$), there is a positive constant \bar{M} such that $|F_i[\Phi](z)| \leq \bar{M}$ for all $\Phi \in \Gamma$, $z \in \mathbb{R}$ and $i = 1, 2, 3$, hence $|P_i[\Phi](z)| \leq \bar{M}/\beta$ and $|P_i[\Phi]'(z)| = |\beta P_i[\Phi](z) - F_i[\Phi](z)|/s \leq 2\bar{M}/s$ for all $\Phi \in \Gamma$, $z \in \mathbb{R}$ and $i = 1, 2, 3$. Therefore, from Arzelà-Ascoli theorem (applied in each compact interval $[-m, m]$ with $m \in \mathbb{N}$), together with a diagonal extraction process and the

definition (2.9) of the norm $\|\cdot\|_X$ in X , it follows that each sequence in $P(\Gamma)$ has a converging subsequence in $(X, \|\cdot\|_X)$ (the limit then belongs to Γ from the previous paragraph and the closedness of Γ).

Furthermore, for any $\Phi \in \Gamma$ and any sequence $(\Phi_n)_{n \in \mathbb{N}}$ in Γ such that $\|\Phi_n - \Phi\|_X \rightarrow 0$ as $n \rightarrow +\infty$, one has in particular $\Phi_n(z) \rightarrow \Phi(z)$ as $n \rightarrow +\infty$ in \mathbb{R}^3 for every $z \in \mathbb{R}$ and, together with the boundedness of the sequence $(\Phi_n)_{n \in \mathbb{N}}$ in $L^\infty(\mathbb{R})^3$ and the Lebesgue dominated convergence theorem, one gets that $F_i[\Phi_n](z) \rightarrow F_i[\Phi](z)$ as $n \rightarrow +\infty$ for every $z \in \mathbb{R}$ and $i = 1, 2, 3$. With the boundedness of the sequence $(F_1[\Phi_n], F_2[\Phi_n], F_3[\Phi_n])_{n \in \mathbb{N}}$ in $L^\infty(\mathbb{R})^3$, one infers that $P[\Phi_n](z) \rightarrow P[\Phi](z)$ as $n \rightarrow +\infty$ in \mathbb{R}^3 for every $z \in \mathbb{R}$. Therefore, each converging subsequence of the sequence $(P[\Phi_n])_{n \in \mathbb{N}}$ in $(X, \|\cdot\|_X)$ must converge to the unique possible limit $P[\Phi]$, and finally $P[\Phi_n] \rightarrow P[\Phi]$ in X as $n \rightarrow +\infty$.

We conclude from the previous two paragraphs that $P : \Gamma \rightarrow \Gamma$ is completely continuous. It thus follows from the Schauder fixed-point theorem that P has a fixed-point $\Phi = (\phi_1, \phi_2, \phi_3)$ in Γ , which is then a $C^1(\mathbb{R})$ solution of (1.3) such that $\underline{\phi}_i \leq \phi_i \leq \bar{\phi}_i$ in \mathbb{R} for $i = 1, 2, 3$. This completes the proof of the lemma. \square

Moreover, due to the negativity of $\alpha(-\infty)$, we have the following universal result for the left-hand tail limit. Since its proof is almost the same as that of [10, (3.19)] (see also [32]), we omit it here safely.

Proposition 2.3. *It holds $(\phi_1, \phi_2, \phi_3)(-\infty) = (0, 0, 0)$ for any nonnegative bounded solution (ϕ_1, ϕ_2, ϕ_3) of (1.3).*

Note that any nonnegative nontrivial (in the sense that each component is not identically 0 in \mathbb{R}) solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) must be positive (in the sense that $\phi_i > 0$ in \mathbb{R} for $i = 1, 2, 3$), by a property similar to the strong maximum principle: indeed, if $\phi_i(z_0) = 0$ for some $i \in \{1, 2, 3\}$ and $z_0 \in \mathbb{R}$, then $\phi'_i(z_0) = 0$, and $(J_i * \phi_i)(z_0) = 0$ from the equation, hence $\phi_i = 0$ in $[z_0 - y_i^\pm - \eta_i, z_0 - y_i^\pm + \eta_i]$ with $\eta_i > 0$ and $-\infty < y_i^- < 0 < y_i^+ < +\infty$ as in (J1); one would then get that $\phi_i = 0$ in $[z_0 - ky_i^- - k\eta_i, z_0 - ky_i^- + k\eta_i]$ for all $k \in \mathbb{N}$ by an immediate induction; by choosing positive integers n and k such that $z_0 + ny_i^+ \in [z_0 - ky_i^- - k\eta_i, z_0 - ky_i^- + k\eta_i]$, it follows that $\phi_i(z_0 + ny_i^+) = 0$, and then $\phi_i = 0$ in $[z_0 + (n-1)y_i^+ - \eta_i, z_0 + (n-1)y_i^+ + \eta_i]$ and also $\phi_i = 0$ in $[z_0 - n\eta_i, z_0 + n\eta_i]$ by induction; finally the non-empty set $\{z \in \mathbb{R} : \phi_i(z) = 0\}$ would be both open and closed (by continuity of ϕ_i), hence $\phi_i \equiv 0$ in \mathbb{R} , a contradiction.

It is interesting to note from the boundedness of α (since it is continuous and has finite limits at $\pm\infty$) and $(\alpha 2)$ that $\alpha(+\infty) - \alpha(z) \leq Ce^{-\rho z}$ for all $z \in \mathbb{R}$, if we choose the constant C larger. Then for any positive constant A it holds

$$\alpha(z + A) \geq \alpha(+\infty) - \varepsilon e^{-\rho z} \text{ for all } z \in \mathbb{R}, \text{ with } \varepsilon := Ce^{-\rho A}.$$

Observe that $\varepsilon > 0$ can be made as small as we want, if we choose A large enough. We further note that $(\phi_1, \phi_2, \phi_3)(\cdot + A)$ is a solution of (1.3) if and only if (ϕ_1, ϕ_2, ϕ_3) is a solution of (1.3) with α replaced by $\alpha(\cdot + A)$. Therefore, in the sequel, (up to a translation) condition $(\alpha 2)$ can be rephrased without loss of generality as

$$(2.10) \quad \alpha(z) \geq \alpha(+\infty) - \varepsilon e^{-\rho z} \text{ for all } z \in \mathbb{R},$$

for a suitable choice of $\varepsilon > 0$ as small as we need.

Now, we consider the following scalar equation

$$(2.11) \quad -s\phi'(z) = d\mathcal{N}[\phi](z) + r\phi(z)[\alpha(z) - \phi(z)], \quad z \in \mathbb{R},$$

where parameters s, d, r are positive constants, and

$$\mathcal{N}[\phi](z) := \int_{\mathbb{R}} J(y)\phi(z-y)dy - \phi(z)$$

in which the kernel J is assumed to satisfy (J1)-(J3) with some constants $\eta > 0$, $-\infty < y^- < 0 < y^+ < +\infty$ in (J1), and $-\infty \leq \tilde{\lambda} < 0 < \hat{\lambda} \leq +\infty$ in (J3), and the continuous heterogeneous function α is assumed to satisfy $(\alpha 1)$ - $(\alpha 2)$, but its limit at $+\infty$ is not assumed to be equal to 1 in general, that is, (1.4) is not assumed here. Recall from [27, Theorem 4.5] that a positive nondecreasing solution ϕ of (2.11) satisfying

$$(2.12) \quad \phi(-\infty) = 0 \quad \text{and} \quad \phi(+\infty) = \alpha(+\infty),$$

is known to exist when J is assumed to be even and compactly supported and when α is assumed to be nondecreasing. Without the symmetry of J and the assumption of compact support of J , and without the monotonicity of α , we still get the existence of a forced wave for (2.11)-(2.12) (which may nevertheless not be monotone in general):

Proposition 2.4. *Under assumptions (J1)-(J3) on J , with some constants $\eta > 0$, $-\infty < y^- < 0 < y^+ < +\infty$ in (J1), and $-\infty \leq \tilde{\lambda} < 0 < \hat{\lambda} \leq +\infty$ in (J3), and under conditions $(\alpha 1)$ - $(\alpha 2)$ on α , problem (2.11)-(2.12) with $s > 0$ admits a positive $C^1(\mathbb{R})$ solution ϕ such that $\alpha(+\infty) \geq \phi(z) \geq \alpha(+\infty) - Be^{-\lambda_0 z}$ for all $z \in \mathbb{R}$, for some $\lambda_0 > 0$ and $B > 0$.*

Proof. First, we give a pair of super-sub-solutions to (2.11) as follows. It is clear that the constant function $\bar{\phi}(z) \equiv \alpha(+\infty)$ is a super-solution of (2.11), in the sense that

$$d\mathcal{N}[\bar{\phi}](z) + s\bar{\phi}'(z) + r\bar{\phi}(z)[\alpha(z) - \bar{\phi}(z)] \leq 0 \quad \text{for all } z \in \mathbb{R}.$$

For the sub-solution, we consider the nonnegative bounded function

$$\underline{\phi}(z) := \max\{\alpha(+\infty) - e^{-\lambda_0 z}, 0\}$$

for some small positive constant $\lambda_0 \in (0, \rho)$, where $\rho > 0$ is the positive constant appearing in $(\alpha 2)$. To choose λ_0 , we let

$$g(\lambda) := d[I(\lambda) - 1] - s\lambda, \quad I(\lambda) := \int_{\mathbb{R}} J(y)e^{\lambda y} dy, \quad \lambda \in [0, \hat{\lambda}).$$

Note that $g(0) = 0$ and $g'(0) = -s$, since

$$g'(\lambda) = d \int_{\mathbb{R}} yJ(y)e^{\lambda y} dy - s \quad \text{and} \quad \int_{\mathbb{R}} yJ(y)dy = 0$$

due to the assumptions (J1)-(J3). Hence there is a $\lambda_0 \in (0, \min\{\rho, \hat{\lambda}\})$ such that $g(\lambda_0) < 0$. Now, we choose $\varepsilon > 0$ small enough such that

$$(2.13) \quad g(\lambda_0) + r\varepsilon\alpha(+\infty)^{\rho/\lambda_0} = g(\lambda_0) + r\varepsilon e^{(\rho/\lambda_0)\ln\alpha(+\infty)} < 0$$

and (2.10) holds for this ε . Then for $z > -\lambda_0^{-1} \ln \alpha(+\infty) =: z_0$ we have $\underline{\phi}(z) = \alpha(+\infty) - e^{-\lambda_0 z} > 0$ and, since $\underline{\phi}(x) \geq \alpha(+\infty) - e^{-\lambda_0 x}$ for all $x \in \mathbb{R}$, we get that

$$\begin{aligned} & d\mathcal{N}[\underline{\phi}](z) + s\underline{\phi}'(z) + r\underline{\phi}(z)[\alpha(z) - \underline{\phi}(z)] \\ & \geq d \int_{\mathbb{R}} J(y)(\alpha(+\infty) - e^{-\lambda_0(z-y)}) dy - d(\alpha(+\infty) - e^{-\lambda_0 z}) + s\lambda_0 e^{-\lambda_0 z} + r\underline{\phi}(z)[\alpha(z) - \underline{\phi}(z)] \\ & = -e^{-\lambda_0 z} \{d[I(\lambda_0) - 1] - s\lambda_0\} + r(\alpha(+\infty) - e^{-\lambda_0 z})(\alpha(z) - \alpha(+\infty) + e^{-\lambda_0 z}) \\ & \geq -e^{-\lambda_0 z} \{d[I(\lambda_0) - 1] - s\lambda_0\} - r\varepsilon(\alpha(+\infty) - e^{-\lambda_0 z})e^{-\rho z} \\ & \geq -e^{-\lambda_0 z} \{d[I(\lambda_0) - 1] - s\lambda_0 + r\varepsilon\alpha(+\infty)e^{-(\rho-\lambda_0)z}\} \geq -e^{-\lambda_0 z} [g(\lambda_0) + r\varepsilon e^{(\rho/\lambda_0) \ln \alpha(+\infty)}] > 0, \end{aligned}$$

using (2.10) and (2.13), together with $\lambda_0 < \rho$ and $z > z_0 = -\lambda_0^{-1} \ln \alpha(+\infty)$. Furthermore, for $z < z_0$, we have $\underline{\phi} = 0$ in a neighborhood of z and $d\mathcal{N}[\underline{\phi}](z) + s\underline{\phi}'(z) + r\underline{\phi}(z)[\alpha(z) - \underline{\phi}(z)] = d(J * \underline{\phi})(z) \geq 0$. Hence $\underline{\phi}$ is a sub-solution of (2.11), in a sense similar to Definition 2.1, that is, it is continuous in \mathbb{R} , of class C^1 in $\mathbb{R} \setminus \{z_0\}$, and it satisfies $d\mathcal{N}[\underline{\phi}](z) + s\underline{\phi}'(z) + r\underline{\phi}(z)[\alpha(z) - \underline{\phi}(z)] \geq 0$ for all $z \in \mathbb{R} \setminus \{z_0\}$.

Next, since $0 \leq \underline{\phi} \leq \bar{\phi} = \alpha(+\infty)$, it follows with the same arguments as in the proof of Lemma 2.2 that there is a $C^1(\mathbb{R})$ solution ϕ of (2.11) such that $0 \leq \underline{\phi} \leq \phi \leq \bar{\phi} = \alpha(+\infty)$ in \mathbb{R} . Hence, $\phi(+\infty) = \alpha(+\infty)$, since $\underline{\phi}(+\infty) = \bar{\phi}(+\infty) = \alpha(+\infty)$. Furthermore, from the definition of $\underline{\phi}$ and the boundedness of $\bar{\phi}$, there is $B \geq 1$ such that $\phi(z) \geq \alpha(+\infty) - Be^{-\lambda_0 z}$ for all $z \in \mathbb{R}$. Finally, ϕ is positive due to the strong maximum principle (as for the proof of the positivity of (ϕ_1, ϕ_2, ϕ_3) after the statement of Proposition 2.3) and the property $\phi(-\infty) = 0$ can be proved in the same way as that in [10, (3.19)] (see also [32]), since $\alpha(-\infty) < 0$. We omit it here. This completes the proof of the proposition. \square

We remark here that the solution ϕ of (2.11)-(2.12) obtained in Proposition 2.4 also satisfies condition of the type (a2) by construction, since it is trapped between $\underline{\phi}(z) = \max\{\phi(+\infty) - e^{-\lambda_0 z}, 0\}$ and $\bar{\phi}(z) = \phi(+\infty)$.

3. PROOF OF THEOREM 1.1

Throughout this section, we suppose (1.2) and (1.4)-(1.5). The proof of Theorem 1.1 is similar to that of [10, Theorem 2.1]. We only provide here a brief outline of the proof as follows. We fix any $s > 0$. Recall from (1.2) and (1.5) that

$$\gamma_3 := 1 - 2b(a - 1) > 0, \quad \gamma_2 := (1 - h - 2ab)(a - 1) > 0, \quad \gamma_1 := (1 - k - 2ab)(a - 1) > 0.$$

First, let $\underline{\phi}_3$ be a positive $C^1(\mathbb{R})$ solution of

$$\begin{cases} -s\underline{\phi}_3'(z) = d_3\mathcal{N}_3[\underline{\phi}_3](z) + r_3\underline{\phi}_3(z)[\alpha(z) - 2b(a - 1) - \underline{\phi}_3(z)], & z \in \mathbb{R}, \\ \underline{\phi}_3(-\infty) = 0, \quad \underline{\phi}_3(+\infty) = \gamma_3. \end{cases}$$

Such $\underline{\phi}_3$ exists, by Proposition 2.4 applied with the continuous function $\alpha - 2b(a - 1)$ instead of α , with limit $\gamma_3 = \alpha(+\infty) - 2b(a - 1) = 1 - 2b(a - 1) > 0$ at $+\infty$ and limit $\alpha(-\infty) - 2b(a - 1) < 0$ at $-\infty$. Furthermore, by Proposition 2.4, $\underline{\phi}_3$ is constructed such that $\underline{\phi}_3 \leq \phi_3(+\infty) = \gamma_3$ in \mathbb{R} , and $\underline{\phi}_3$ converges to γ_3 exponentially at $+\infty$.

Next, note that the continuous bounded function $a\underline{\phi}_3 - 1 - h(a - 1)$ is such that

$$a\underline{\phi}_3(-\infty) - 1 - h(a - 1) = -1 - h(a - 1) < 0, \quad a\underline{\phi}_3(+\infty) - 1 - h(a - 1) = \gamma_2 > 0,$$

and $a\underline{\phi}_3 - 1 - h(a - 1) \leq a\underline{\phi}_3(+\infty) - 1 - h(a - 1)$ in \mathbb{R} . By Proposition 2.4 again, there exists a positive $C^1(\mathbb{R})$ solution $\underline{\phi}_2$ of

$$\begin{cases} -s\underline{\phi}'_2(z) = d_2\mathcal{N}_2[\underline{\phi}_2](z) + r_2\underline{\phi}_2(z)[a\underline{\phi}_3(z) - 1 - h(a - 1) - \underline{\phi}_2(z)], & z \in \mathbb{R}, \\ \underline{\phi}_2(-\infty) = 0, \quad \underline{\phi}_2(+\infty) = \gamma_2, \end{cases}$$

such that $\underline{\phi}_2 \leq \underline{\phi}_2(+\infty) = \gamma_2$ in \mathbb{R} and $\underline{\phi}_2$ converges to γ_2 exponentially at $+\infty$. Similarly, there exists a positive $C^1(\mathbb{R})$ solution $\underline{\phi}_1$ of

$$\begin{cases} -s\underline{\phi}'_1(z) = d_1\mathcal{N}_1[\underline{\phi}_1](z) + r_1\underline{\phi}_1(z)[a\underline{\phi}_3(z) - 1 - k(a - 1) - \underline{\phi}_1(z)], & z \in \mathbb{R}, \\ \underline{\phi}_1(-\infty) = 0, \quad \underline{\phi}_1(+\infty) = \gamma_1 > 0, \end{cases}$$

such that $\underline{\phi}_1 \leq \underline{\phi}_1(+\infty) = \gamma_1$ in \mathbb{R} and $\underline{\phi}_1$ converges to γ_1 exponentially at $+\infty$.

It is straightforward to verify that $(\bar{\phi}_1, \bar{\phi}_2, \bar{\phi}_3) \equiv (a - 1, a - 1, 1)$ and $(\underline{\phi}_1, \underline{\phi}_2, \underline{\phi}_3)$ are a pair of $C^1(\mathbb{R})$ nonnegative bounded upper-lower-solutions of (1.3), such that $\underline{\phi}_i \leq \bar{\phi}_i$ in \mathbb{R} for all $i = 1, 2, 3$ (the last property holds since $(\gamma_1, \gamma_2, \gamma_3) < (a - 1, a - 1, 1)$ componentwise). Hence the existence of positive wave profile (ϕ_1, ϕ_2, ϕ_3) of (1.3) follows from Lemma 2.2, with

$$0 < \underline{\phi}_i \leq \phi_i \leq \bar{\phi}_i \quad \text{in } \mathbb{R} \text{ for } i = 1, 2, 3.$$

Note that

$$\phi_i^- := \liminf_{z \rightarrow +\infty} \phi_i(z) \geq \gamma_i > 0, \quad i = 1, 2, 3.$$

With this information, the same proof of [10, Lemma 3.4] can be applied to derive that $(\phi_1, \phi_2, \phi_3)(+\infty) = E_4$. We also refer the reader to the proof of [21, (4.1)]. Indeed, setting

$$\phi_i^+ := \limsup_{z \rightarrow +\infty} \phi_i(z), \quad i = 1, 2, 3,$$

picking any $\varepsilon > 0$ small enough and considering the family of contracting parallelepipeds $\prod_{i=1}^3 [m_i(\theta), M_i(\theta)]$ (with $\theta \in [0, 1]$) defined by

$$\begin{cases} m_1(\theta) := \theta u^* + (1 - \theta)(\gamma_1 - \varepsilon), & M_1(\theta) := \theta u^* + (1 - \theta)(a - 1 + \varepsilon), \\ m_2(\theta) := \theta v^* + (1 - \theta)(\gamma_2 - \varepsilon), & M_2(\theta) := \theta v^* + (1 - \theta)(a - 1 + \varepsilon), \\ m_3(\theta) := \theta w^* + (1 - \theta)(\gamma_3 - \varepsilon^2), & M_3(\theta) := \theta u^* + (1 - \theta)(1 + \varepsilon^2), \end{cases}$$

we can show that $0 \in \Theta$ and $\sup \Theta = 1$, where

$$\Theta := \{\theta \in [0, 1] \mid m_i(\theta) < \phi_i^- \leq \phi_i^+ < M_i(\theta), \quad i = 1, 2, 3\}.$$

The details can be found in [10, 21]. We omit them here. Hence $(\phi_1, \phi_2, \phi_3)(+\infty) = (u^*, v^*, w^*) = E_4$, and the proof of Theorem 1.1 is complete, thanks to Proposition 2.3.

4. PROOF OF THEOREM 1.2

Throughout this section, we suppose (1.2) and (1.4). For the sufficiency part, let us assume without loss of generality that $s_1^* \geq s_2^*$, with s_1^* and s_2^* defined by (1.6), and let us then fix $s > s_1^*$. First, we set

$$(4.1) \quad g_i(\lambda) := d_i[I_i(\lambda) - 1] - s\lambda + r_i(a - 1), \quad i = 1, 2, \quad \lambda \in [0, \hat{\lambda}_i).$$

From (1.2) and (J1)-(J3) (and the comments after (J1)-(J3)), there exist $0 < \lambda_1 < \lambda_2 < \hat{\lambda}_1$ such that $g_1(\lambda_1) = g_1(\lambda_2) = 0$, $g_1(\lambda) < 0$ if and only if $\lambda \in (\lambda_1, \lambda_2)$, and $g_1(\lambda) \leq 0$ if and only if $\lambda \in [\lambda_1, \lambda_2]$. Similarly, since $s > s_2^*$, the equation $g_2(\lambda) = 0$ has two positive roots λ_3, λ_4 with $0 < \lambda_3 < \lambda_4 < \hat{\lambda}_2$. Moreover, $g_2(\lambda) < 0$ if and only if $\lambda \in (\lambda_3, \lambda_4)$, and $g_2(\lambda) \leq 0$ if and only if $\lambda \in [\lambda_3, \lambda_4]$. Set now

$$(4.2) \quad g_3(\lambda) := d_3[I_3(\lambda) - 1] - s\lambda.$$

Note that $g_3(0) = 0$ and $g_3'(0) = -s < 0$. Hence we can choose

$$(4.3) \quad 0 < \lambda_0 < \min\{\lambda_1, \lambda_3, \rho, \hat{\lambda}_3\}$$

such that

$$g_3(\lambda_0) < 0,$$

where $\rho > 0$ is as in (α2). We then define, for $z \in \mathbb{R}$,

$$(4.4) \quad \begin{cases} \bar{\phi}_1(z) = \min\{(a-1)e^{-\lambda_1 z}, a-1\}, & \underline{\phi}_1(z) = \max\{(a-1)e^{-\lambda_1 z} - p_1 e^{-\mu_1 z}, 0\}, \\ \bar{\phi}_2(z) = \min\{(a-1)e^{-\lambda_3 z}, a-1\}, & \underline{\phi}_2(z) = \max\{(a-1)e^{-\lambda_3 z} - p_2 e^{-\mu_2 z}, 0\}, \\ \bar{\phi}_3(z) = 1, & \underline{\phi}_3(z) = \max\{1 - e^{-\lambda_0 z}, 0\}, \end{cases}$$

where the positive constants $\mu_i, p_i, i = 1, 2$, are chosen (in the following order) such that

$$(4.5) \quad \lambda_1 < \mu_1 < \min\{\lambda_2, \lambda_1 + \lambda_0\},$$

$$(4.6) \quad p_1 > \max\left\{a-1, \frac{r_1(a-1)[2a-1+k(a-1)]}{-g_1(\mu_1)}\right\},$$

$$(4.7) \quad \lambda_3 < \mu_2 < \min\{\lambda_4, \lambda_3 + \lambda_0\},$$

$$(4.8) \quad p_2 > \max\left\{a-1, \frac{r_2(a-1)[2a-1+h(a-1)]}{-g_2(\mu_2)}\right\}.$$

Note that the continuous functions $\underline{\phi}_i$ and $\bar{\phi}_i$ are nonnegative, bounded in \mathbb{R} , of class C^1 except at finitely many real numbers, and that $\underline{\phi}_i \leq \bar{\phi}_i$ in \mathbb{R} , for $i = 1, 2, 3$.

Then we have

Lemma 4.1. *With $s > s_1^* \geq s_2^*$ and the above notations, there exists a positive solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) such that $\underline{\phi}_i \leq \phi_i \leq \bar{\phi}_i$ in \mathbb{R} , $i = 1, 2, 3$.*

Proof. We choose $\varepsilon > 0$ small enough such that

$$(4.9) \quad g_3(\lambda_0) + \varepsilon r_3 < 0.$$

Also, up to a translation, condition (2.10) holds for this ε .

We now verify (2.1)-(2.6) hold for $(\bar{\phi}_1, \bar{\phi}_2, \bar{\phi}_3)$ and $(\underline{\phi}_1, \underline{\phi}_2, \underline{\phi}_3)$ defined in (4.4) for all $z \in \mathbb{R}$ except finitely many points. For (2.1), the inequality is obvious for $z < 0$ since $\bar{\phi}_1 \leq a - 1$ in \mathbb{R} , $\bar{\phi}_1 = a - 1$ in $(-\infty, 0]$, and $\bar{\phi}_3 \equiv 1$ in \mathbb{R} . It then suffices to consider $z > 0$. As in the proof of Proposition 2.4, using $\bar{\phi}_1(y) \leq (a - 1)e^{-\lambda_1 y}$ for all $y \in \mathbb{R}$ and $\bar{\phi}_1(z) = (a - 1)e^{-\lambda_1 z}$ for all $z > 0$, we have

$$(4.10) \quad d_1 \mathcal{N}_1[\bar{\phi}_1](z) + s\bar{\phi}_1'(z) \leq (a - 1)e^{-\lambda_1 z} \{d_1[I_1(\lambda_1) - 1] - s\lambda_1\} \text{ for all } z > 0,$$

and thus

$$(4.11) \quad \begin{aligned} & d_1 \mathcal{N}_1[\bar{\phi}_1](z) + s\bar{\phi}_1'(z) + r_1 \bar{\phi}_1(z) [-1 - \bar{\phi}_1(z) - k\underline{\phi}_2(z) + a\bar{\phi}_3(z)] \\ & \leq (a - 1)e^{-\lambda_1 z} \{d_1[I_1(\lambda_1) - 1] - s\lambda_1 + r_1(-1 + a)\} = 0. \end{aligned}$$

Hence (2.1) holds for all $z \neq 0$. Similarly, (2.2) holds for all $z \neq 0$, using

$$(4.12) \quad g_2(\lambda_3) = d_2[I_2(\lambda_3) - 1] - s\lambda_3 + r_2(a - 1) = 0.$$

It is trivial that (2.3) holds for all $z \in \mathbb{R}$, since $\alpha(z) \leq \alpha(+\infty) = 1$ and $\underline{\phi}_1(z) \geq 0$, $\underline{\phi}_2(z) \geq 0$ for all $z \in \mathbb{R}$.

For (2.4), since $p_1 > a - 1 > 0$ and $\mu_1 > \lambda_1 > 0$ there is a unique $z_1 := (\mu_1 - \lambda_1)^{-1} \ln(p_1 / (a - 1)) > 0$ such that $\underline{\phi}_1(z) = 0$ for all $z \leq z_1$ and

$$\underline{\phi}_1(z) = (a - 1)e^{-\lambda_1 z} - p_1 e^{-\mu_1 z} > 0 \text{ for all } z > z_1.$$

It is trivial that (2.4) holds for $z < z_1$. On the other hand, since $\underline{\phi}_1(y) \geq (a - 1)e^{-\lambda_1 y} - p_1 e^{-\mu_1 y}$ for all $y \in \mathbb{R}$ with equality in $[z_1, +\infty)$, with $0 < \lambda_1 < \mu_1 < \lambda_2 < \hat{\lambda}_1$, it follows that, for every $z > z_1$,

$$(4.13) \quad d_1 \mathcal{N}_1[\underline{\phi}_1](z) + s\underline{\phi}_1'(z) \geq (a - 1)e^{-\lambda_1 z} \{d_1[I_1(\lambda_1) - 1] - s\lambda_1\} - p_1 e^{-\mu_1 z} \{d_1[I_1(\mu_1) - 1] - s\mu_1\},$$

and

$$\begin{aligned} & r_1 \underline{\phi}_1(z) [-1 - \underline{\phi}_1(z) - k\bar{\phi}_2(z) + a\underline{\phi}_3(z)] \\ & \geq r_1 [(a - 1)e^{-\lambda_1 z} - p_1 e^{-\mu_1 z}] \{-1 - (a - 1)e^{-\lambda_1 z} - k(a - 1)e^{-\lambda_3 z} + a - a e^{-\lambda_0 z}\} \\ & \geq r_1 (a - 1) [(a - 1)e^{-\lambda_1 z} - p_1 e^{-\mu_1 z}] - r_1 (a - 1) e^{-\lambda_1 z} [(a - 1)(e^{-\lambda_1 z} + k e^{-\lambda_3 z}) + a e^{-\lambda_0 z}]. \end{aligned}$$

Hence, since $g_1(\lambda_1) = 0$, we have, for $z > z_1$,

$$\begin{aligned} \mathcal{L}_1(z) & \geq e^{-\mu_1 z} \{-p_1 g_1(\mu_1) - r_1 (a - 1) [(a - 1)(e^{(\mu_1 - 2\lambda_1)z} + k e^{(\mu_1 - \lambda_1 - \lambda_3)z}) + a e^{(\mu_1 - \lambda_1 - \lambda_0)z}]\} \\ & \geq e^{-\mu_1 z} \{-p_1 g_1(\mu_1) - r_1 (a - 1) [2a - 1 + k(a - 1)]\} > 0, \end{aligned}$$

due to (4.3) and (4.5)-(4.6). Hence (2.4) holds for all $z \neq z_1$.

Similarly, since $p_2 > a - 1 > 0$ and $\mu_2 > \lambda_3 > 0$, there is a unique $z_2 := (\mu_2 - \lambda_3)^{-1} \ln(p_2 / (a - 1)) > 0$ such that $\underline{\phi}_2(z) = 0$ for all $z \leq z_2$ and

$$\underline{\phi}_2(z) = (a - 1)e^{-\lambda_3 z} - p_2 e^{-\mu_2 z} > 0 \text{ for all } z > z_2.$$

Then, as in the previous paragraph, we can easily check that (2.5) holds for all $z \neq z_2$, using (4.3) and (4.7)-(4.8), together with $g_2(\lambda_3) = 0$. We omit the details.

Finally, (2.6) holds trivially for $z < 0$. For $z > 0$, we compute, using (1.4) and (2.10),

$$\begin{aligned} \mathcal{L}_3(z) &= d_3 \mathcal{N}_3[\underline{\phi}_3](z) + s \underline{\phi}_3'(z) + r_3 \underline{\phi}_3(z) [\alpha(z) - b \bar{\phi}_1(z) - b \bar{\phi}_2(z) - \underline{\phi}_3(z)] \\ &\geq -e^{-\lambda_0 z} \{d_3 [I_3(\lambda_0) - 1] - s \lambda_0\} \\ &\quad + r_3 (1 - e^{-\lambda_0 z}) [1 - \varepsilon e^{-\rho z} - b(a-1)e^{-\lambda_1 z} - b(a-1)e^{-\lambda_3 z} - 1 + e^{-\lambda_0 z}] \\ &\geq -e^{-\lambda_0 z} \{d_3 [I_3(\lambda_0) - 1] - s \lambda_0\} - \varepsilon r_3 (1 - e^{-\lambda_0 z}) e^{-\rho z}, \end{aligned}$$

using $2b(a-1) < 1$ (by (1.2)) and $e^{-\lambda_0 z} \geq e^{-\lambda_j z}$ for $z > 0$ (since $\lambda_0 < \lambda_j$) for $j = 1, 3$. Moreover, since $0 < \lambda_0 < \rho$ by (4.3), we deduce that

$$\mathcal{L}_3(z) \geq -e^{-\lambda_0 z} \{d_3 [I_3(\lambda_0) - 1] - s \lambda_0 + r_3 \varepsilon\} > 0 \text{ for all } z > 0,$$

by the choice of ε in (4.9). Hence (2.6) holds for all $z \neq 0$.

Lemma 2.2 then yields the existence of a nonnegative bounded solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) such that $\underline{\phi}_i \leq \phi_i \leq \bar{\phi}_i$ in \mathbb{R} for $i = 1, 2, 3$. In particular, each function ϕ_i is nonnegative and nontrivial, hence it is positive in \mathbb{R} by the strong maximum principle, as explained after Proposition 2.3. The proof of Lemma 4.1 is thereby complete. \square

Proof of Theorem 1.2. By Lemma 4.1 and the definition of the upper-lower-solutions in (4.4), the positive solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) given in Lemma 4.1 satisfies

$$(\phi_1, \phi_2, \phi_3)(+\infty) = (0, 0, 1) = E_1.$$

Then, together with Proposition 2.3, the existence part of Theorem 1.2 follows.

For the only if part, we follow the idea of [10, Theorem 3.5]. Assume that there exists a positive solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) and (1.7) for some $s > 0$, and assume that $(\tilde{\lambda}_i, \hat{\lambda}_i) = (-\infty, +\infty)$ for some $i \in \{1, 2\}$. Without loss of generality, let us assume that $i = 1$. Set $\zeta(z) := \phi_1'(z)/\phi_1(z)$. Then ζ satisfies

$$-s\zeta(z) = d_1 \left[\int_{\mathbb{R}} J_1(y) e^{\int_z^{z-y} \zeta(x) dx} dy - 1 \right] + r_1 [-1 - \phi_1(z) - k\phi_2(z) + a\phi_3(z)], \quad z \in \mathbb{R}.$$

It follows from [36, Proposition 3.7]² that the limit $-\lambda := \lim_{z \rightarrow +\infty} \zeta(z)$ exists in \mathbb{R} (necessarily, $\lambda \geq 0$, since $\phi_1 > 0$ in \mathbb{R} and $\phi_1(+\infty) = 0$) and λ satisfies

$$s\lambda = d_1 [I_1(\lambda) - 1] + r_1 (a - 1).$$

Since $I_1(0) = 1$ and $r_1(a-1) > 0$, it follows that $\lambda > 0$, hence we obtain that $s \geq s_1^*$, owing to the definition of s_1^* in (1.6). Thereby the proof is complete. \square

5. PROOF OF THEOREM 1.3

Throughout this section, we suppose (1.2) and (1.4), together with (1.10)-(1.11). For the sufficiency part, let us assume that $s \geq R_2(\rho)$ and $s > s_2^{**}$, where $\rho > 0$ is the constant

²Notice that, in addition to the assumption $(\tilde{\lambda}_1, \hat{\lambda}_1) = (-\infty, +\infty)$, J_1 is assumed to be even and of class $C^1(\mathbb{R})$ in [36], but the proof of [36, Proposition 3.7] still works under assumptions (J1)-(J3) for J_1 , under the assumption $(\tilde{\lambda}_1, \hat{\lambda}_1) = (-\infty, +\infty)$.

in (α2), and R_2 and s_2^{**} are defined in (1.9). We recall that $\beta_2 > 0$ is defined in (1.8). From (J1)-(J3) and (1.2), the function G_2 defined by

$$(5.1) \quad G_2(\lambda) := d_2[I_2(\lambda) - 1] - s\lambda + r_2\beta_2, \quad \lambda \in [0, \hat{\lambda}_2),$$

has two positive zeroes λ_5, λ_6 such that $0 < \lambda_5 < \lambda_6 < \hat{\lambda}_2$. Note that $G_2(\lambda) < 0$ if and only if $\lambda \in (\lambda_5, \lambda_6)$, and $G_2(\lambda) \leq 0$ if and only if $\lambda \in [\lambda_5, \lambda_6]$. Since $s \geq R_2(\rho)$ and $\rho > 0$, one necessarily has $\rho < \hat{\lambda}_2$ and $G_2(\rho) \leq 0$, hence $\rho \geq \lambda_5$.

We recall that $u_p = (a - 1)/(ab + 1)$ and $w_p = (b + 1)/(ab + 1)$ and we define, for $z \in \mathbb{R}$,

$$(5.2) \quad \begin{cases} \bar{\phi}_1(z) := \min\{u_p + Ae^{-\lambda_5 z}, a - 1\}, & \underline{\phi}_1(z) := \max\{u_p(1 - e^{-\lambda_5 z}), 0\}, \\ \bar{\phi}_2(z) := \min\{(a - 1)e^{-\lambda_5 z}, a - 1\}, & \underline{\phi}_2(z) := \max\{(a - 1)e^{-\lambda_5 z} - qe^{-\nu z}, 0\}, \\ \bar{\phi}_3(z) := \min\{w_p + bu_p e^{-\lambda_5 z}, 1\}, & \underline{\phi}_3(z) := \max\{w_p(1 - e^{-\lambda_5 z}), 0\}, \end{cases}$$

where

$$(5.3) \quad A := a - 1 - u_p$$

and the positive constants ν and q are chosen to satisfy

$$(5.4) \quad \lambda_5 < \nu < \min\{\lambda_6, 2\lambda_5\}, \quad q > \max\left\{a - 1, \frac{r_2(a - 1)[(1 + h)(a - 1) + aw_p]}{-G_2(\nu)}\right\}.$$

Note that $0 < u_p < a - 1$ (hence, $A > 0$), $0 < w_p < 1$, $w_p + bu_p = 1$, and that all functions $\underline{\phi}_i$ and $\bar{\phi}_i$ are continuous, nonnegative, bounded in \mathbb{R} , of class C^1 except at finitely many points, with $\underline{\phi}_i \leq \bar{\phi}_i$ in \mathbb{R} for $i = 1, 2, 3$.

Then we have

Lemma 5.1. *With $s > s_2^{**}$, $s \geq R_2(\rho)$ and the above notations, there exists a positive solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) such that $\underline{\phi}_i \leq \phi_i \leq \bar{\phi}_i$ in \mathbb{R} , $i = 1, 2, 3$.*

Proof. Choose $\varepsilon \in (0, r_2\beta_2/r_3)$ so that, up to a translation, (2.10) holds for this ε . We verify (2.1)-(2.6) hold for $(\bar{\phi}_1, \bar{\phi}_2, \bar{\phi}_3)$ and $(\underline{\phi}_1, \underline{\phi}_2, \underline{\phi}_3)$ defined in (5.2) for all $z \in \mathbb{R}$ except finitely many points.

That (2.1)-(2.3) hold for $z < 0$ is trivial. For $z > 0$, since

$$-1 - \bar{\phi}_1(z) - k\underline{\phi}_2(z) + a\bar{\phi}_3(z) \leq -1 - u_p - Ae^{-\lambda_5 z} + aw_p + abu_p e^{-\lambda_5 z} = 0,$$

we obtain

$$\mathcal{U}_1(z) \leq Ae^{-\lambda_5 z} \{d_1[I_1(\lambda_5) - 1] - s\lambda_5\} \leq Ae^{-\lambda_5 z} \{d_2[I_2(\lambda_5) - 1] - s\lambda_5\} < Ae^{-\lambda_5 z} G_2(\lambda_5) = 0,$$

since $I_1(\lambda_5) - 1 \geq 0$ (from the general properties explained after (J1)-(J3)) and since $d_1 \leq d_2$ and $J_1 = J_2$ in \mathbb{R} by (1.10). Hence (2.1) holds for all $z \neq 0$.

For $z > 0$, we compute

$$\begin{aligned} \mathcal{U}_2(z) &\leq (a - 1)e^{-\lambda_5 z} \{d_2[I_2(\lambda_5) - 1] - s\lambda_5\} \\ &\quad + r_2(a - 1)e^{-\lambda_5 z} \{(-1 - hu_p + aw_p) + [hu_p - (a - 1) + abu_p]e^{-\lambda_5 z}\} \\ &= r_2(a - 1)e^{-2\lambda_5 z} [hu_p - (a - 1) + abu_p] \leq 0, \end{aligned}$$

using $-1 - hu_p + aw_p = \beta_2$, $G_2(\lambda_5) = 0$ and $hu_p - (a - 1) + abu_p < (1 + ab)u_p - (a - 1) = 0$. Hence (2.2) holds for all $z \neq 0$.

Still for $z > 0$, we have

$$\alpha(z) - b\underline{\phi}_1(z) - b\underline{\phi}_2(z) - \bar{\phi}_3(z) \leq 1 - bu_p(1 - e^{-\lambda_5 z}) - w_p - bu_p e^{-\lambda_5 z} = 0,$$

hence

$$\mathcal{U}_3(z) \leq bu_p e^{-\lambda_5 z} \{d_3[I_3(\lambda_5) - 1] - s\lambda_5\} \leq bu_p e^{-\lambda_5 z} \{d_2[I_2(\lambda_5) - 1] - s\lambda_5\} < 0,$$

since $I_3(\lambda_5) - 1 \geq 0$, $d_3 \leq d_2$ and $J_3 = J_2$ in \mathbb{R} , by (1.10). Hence (2.3) holds for all $z \neq 0$.

For (2.4), it is trivial for $z < 0$. For $z > 0$, we have

$$d_1 \mathcal{N}_1[\underline{\phi}_1](z) + s\underline{\phi}'_1(z) \geq -u_p e^{-\lambda_5 z} \{d_1[I_1(\lambda_5) - 1] - s\lambda_5\}$$

and

$$r_1 \underline{\phi}_1(z) [-1 - \underline{\phi}_1(z) - k\bar{\phi}_2(z) + a\underline{\phi}_3(z)] = r_1 u_p (1 - e^{-\lambda_5 z}) [-1 - k(a-1)] e^{-\lambda_5 z},$$

using $-1 - u_p + aw_p = 0$. Hence we obtain

$$\begin{aligned} \mathcal{L}_1(z) &\geq -u_p e^{-\lambda_5 z} \{d_1[I_1(\lambda_5) - 1] - s\lambda_5 + r_1[1 + k(a-1)]\} \\ &\geq -u_p e^{-\lambda_5 z} \{d_2[I_2(\lambda_5) - 1] - s\lambda_5 + r_2\beta_2\} = 0, \end{aligned}$$

since $I_1(\lambda_5) - 1 \geq 0$, $d_1 \leq d_2$, $J_1 = J_2$ in \mathbb{R} and $r_1[1 + k(a-1)] \leq r_2\beta_2$, by (1.10)-(1.11). Hence (2.4) holds for all $z \neq 0$.

Note that, since $q > a - 1$ and $\nu > \lambda_5$, there is a unique $z_1 := (\nu - \lambda_5)^{-1} \ln(q/(a-1)) > 0$ such that

$$\underline{\phi}_2(z) = \begin{cases} (a-1)e^{-\lambda_5 z} - qe^{-\nu z} > 0, & z > z_1, \\ 0, & z \leq z_1. \end{cases}$$

Clearly, (2.5) holds for $z < z_1$. For $z > z_1$, we compute

$$d_2 \mathcal{N}_2[\underline{\phi}_2](z) + s\underline{\phi}'_2(z) \geq (a-1)e^{-\lambda_5 z} \{d_2[I_2(\lambda_5) - 1] - s\lambda_5\} - qe^{-\nu z} \{d_2[I_2(\nu) - 1] - s\nu\},$$

using $\underline{\phi}_2(y) \geq (a-1)e^{-\lambda_5 y} - qe^{-\nu y}$ for all $y \in \mathbb{R}$, with equality at z . Also, we have, since $\beta_2 = -1 - hu_p + aw_p$,

$$\begin{aligned} &r_2 \underline{\phi}_2(z) [-1 - h\bar{\phi}_1(z) - \underline{\phi}_2(z) + a\underline{\phi}_3(z)] \\ &= r_2 \underline{\phi}_2(z) \{\beta_2 - h(a-1)e^{-\lambda_5 z} + hu_p e^{-\lambda_5 z} - (a-1)e^{-\lambda_5 z} + qe^{-\nu z} - aw_p e^{-\lambda_5 z}\} \\ &\geq r_2 [(a-1)e^{-\lambda_5 z} - qe^{-\nu z}] \{\beta_2 - [(1+h)(a-1) + aw_p] e^{-\lambda_5 z}\} \\ &\geq r_2 \beta_2 [(a-1)e^{-\lambda_5 z} - qe^{-\nu z}] - r_2 (a-1) [(1+h)(a-1) + aw_p] e^{-2\lambda_5 z}, \end{aligned}$$

for all $z > z_1$. Hence, using $G_2(\lambda_5) = 0$, we obtain, for $z > z_1$,

$$\begin{aligned} \mathcal{L}_2(z) &\geq e^{-\nu z} \{-qG_2(\nu) - r_2(a-1)[(1+h)(a-1) + aw_p] e^{(\nu-2\lambda_5)z}\} \\ &\geq e^{-\nu z} \{-qG_2(\nu) - r_2(a-1)[(1+h)(a-1) + aw_p]\} > 0, \end{aligned}$$

using $\nu < 2\lambda_5$ and (5.4). Hence (2.5) holds for all $z \neq z_1$.

Finally, we consider (2.6). It suffices to consider the case $z > 0$. For $z > 0$, we have $\underline{\phi}_3(z) = w_p - w_p e^{-\lambda_5 z} > 0$, while $\underline{\phi}_3(y) \geq w_p - w_p e^{-\lambda_5 y}$ for all $y \in \mathbb{R}$. Hence

$$\begin{aligned} d_3 \mathcal{N}_3[\underline{\phi}_3](z) + s \underline{\phi}_3'(z) &\geq -w_p e^{-\lambda_5 z} \{d_3 [I_3(\lambda_5) - 1] - s \lambda_5\} \\ &\geq -w_p e^{-\lambda_5 z} \{d_2 [I_2(\lambda_5) - 1] - s \lambda_5\} = r_2 \beta_2 w_p e^{-\lambda_5 z}, \end{aligned}$$

since $I_3(\lambda_5) - 1 \geq 0$, $d_3 \leq d_2$, $J_3 = J_2$ by (1.10), and $G_2(\lambda_5) = 0$. Also, for $z > 0$, using $1 - b u_p - w_p = 0$, $A = a - 1 - u_p$ and $2b(a - 1) < 1$, together with (1.4) and (2.10), we compute

$$\begin{aligned} r_3 \underline{\phi}_3(z) [\alpha(z) - b \bar{\phi}_1(z) - b \bar{\phi}_2(z) - \underline{\phi}_3(z)] &\geq r_3 \underline{\phi}_3(z) \{-\varepsilon e^{-\rho z} + [1 - 2b(a - 1)] e^{-\lambda_5 z}\} \\ &\geq -r_3 w_p \varepsilon e^{-\rho z} \geq -r_3 w_p \varepsilon e^{-\lambda_5 z}, \end{aligned}$$

due to $\rho \geq \lambda_5$. Hence, for $z > 0$, $\mathcal{L}_3(z) \geq r_2 \beta_2 w_p e^{-\lambda_5 z} - r_3 w_p \varepsilon e^{-\lambda_5 z} > 0$ since $\varepsilon < r_2 \beta_2 / r_3$. We conclude that (2.6) holds for all $z \neq 0$.

Since $0 \leq \underline{\phi}_i \leq \bar{\phi}_i$ in \mathbb{R} for $i = 1, 2, 3$, and each function $\underline{\phi}_i$ is nontrivial, the conclusion of Lemma 5.1 follows from Lemma 2.2, as at the end of the proof of Lemma 4.1. \square

With Lemma 5.1, Theorem 1.3 can be proved in the same way as that of Theorem 1.2. We safely omit it here.

6. WAVES WITH CRITICAL SPEED

This section is devoted to the proofs of Theorems 1.4-1.6 on the existence of forced waves with critical speeds. Three cases are considered: two of them are concerned with waves connecting the trivial state $(0, 0, 0)$ and the predator-free state $E_1 = (0, 0, 1)$, and the last one is concerned with waves connecting the trivial state $(0, 0, 0)$ and the mixed state $E_2 = (u_p, 0, w_p)$. Throughout this section, in addition to (1.2) and (1.4), we assume that J_1 is compactly supported. Hence there is a positive constant $S > 0$ such that

$$(6.1) \quad J_1(y) = 0 \quad \text{for almost every } |y| > S.$$

In particular, $\tilde{\lambda}_1 = -\infty$ and $\hat{\lambda}_1 = +\infty$ in (J3).

6.1. Waves connecting $(0, 0, 0)$ and $E_1 = (0, 0, 1)$ in the case $s_1^* = s_2^*$: proof of Theorem 1.4. In this subsection, we assume that $s_1^* = s_2^*$, with $s_i^* > 0$ defined in (1.6), and that J_2 is also compactly supported. Even if it means increasing $S > 0$, we can assume without loss of generality that

$$(6.2) \quad J_2(y) = 0 \quad \text{for almost every } |y| > S.$$

We consider the critical speed

$$s = s_1^* = s_2^*.$$

For $i = 1, 2$, let $\lambda_i^* > 0$ be the unique minimum of Q_i in $(0, \hat{\lambda}_i) = (0, +\infty)$ given by (1.6). With g_i defined in (4.1) with $s = s_1^* = s_2^*$, one has $g_i(\lambda_i^*) = 0$, and λ_i^* is the unique positive root of this equation. Since $Q_i'(\lambda_i^*) = 0$ for $i = 1, 2$, we also have

$$(6.3) \quad s = s_1^* = s_2^* = d_1 \int_{\mathbb{R}} J_1(y) y e^{\lambda_1^* y} dy = d_2 \int_{\mathbb{R}} J_2(y) y e^{\lambda_2^* y} dy.$$

With $\rho > 0$ as in $(\alpha 2)$ and g_3 as in (4.2), we choose a constant $\lambda_0 > 0$ such that

$$0 < \lambda_0 < \min\{\lambda_1^*, \lambda_2^*, \rho\} \quad \text{and} \quad g_3(\lambda_0) = d_3[I_3(\lambda_0) - 1] - s\lambda_0 < 0.$$

We then define

$$(6.4) \quad \begin{cases} \bar{\phi}_1(z) = \begin{cases} (a-1)Bze^{-\lambda_1^*z}, & z > z_1, \\ a-1, & z \leq z_1, \end{cases} & \bar{\phi}_2(z) = \begin{cases} (a-1)Bze^{-\lambda_2^*z}, & z > z_2, \\ a-1, & z \leq z_2, \end{cases} \\ \underline{\phi}_1(z) = \begin{cases} (a-1)Bze^{-\lambda_1^*z} - p_3z^{1/2}e^{-\lambda_1^*z}, & z > z_3, \\ 0, & z \leq z_3, \end{cases} \\ \underline{\phi}_2(z) = \begin{cases} (a-1)Bze^{-\lambda_2^*z} - p_4z^{1/2}e^{-\lambda_2^*z}, & z > z_4, \\ 0, & z \leq z_4, \end{cases} \\ \bar{\phi}_3(z) = 1, \quad \underline{\phi}_3(z) = \max\{1 - e^{-\lambda_0(z-z_0)}, 0\}, \end{cases}$$

where the parameters $B, z_1, z_2, z_0, p_3, z_3, p_4, z_4$ are chosen in the following order:

- $B > \max\{\lambda_1^*e, \lambda_2^*e\}$ large enough such that $z_1 - z'_1 > S$ and $z_2 - z'_2 > S$, where $0 < z'_1 < 1/\lambda_1^* < z_1$, $0 < z'_2 < 1/\lambda_2^* < z_2$, and

$$(6.5) \quad B = \frac{e^{\lambda_1^*z_1}}{z_1} = \frac{e^{\lambda_1^*z'_1}}{z'_1} = \frac{e^{\lambda_2^*z_2}}{z_2} = \frac{e^{\lambda_2^*z'_2}}{z'_2}$$

(notice that the smallest positive root z'_i of each equation $B = e^{\lambda_i^*z}/z$ actually does not appear in (6.4), but it will play a role in the proof of Lemma 6.1 below; observe also that $e^{\lambda_i^*z}/z \leq B$ for $z \in [z'_i, z_i]$, and that $z_i > z'_i + S > S$, for $i = 1, 2$);

- $z_0 > \max\{z_1, z_2\}$ such that

$$(6.6) \quad b(a-1)Bz(e^{-\lambda_1^*z} + e^{-\lambda_2^*z}) \leq e^{-\lambda_0(z-z_0)} \quad \text{for all } z \geq z_0$$

(the choice of z_0 is possible since $0 < \lambda_0 < \min\{\lambda_1^*, \lambda_2^*\}$);

- $p_3 > 0$ large enough such that $z_3 := \{p_3/[(a-1)B]\}^2 > z_0$ and

$$(6.7) \quad p_3 > \frac{8r_1B(a-1) \times \max_{z \geq 0} \{(z+S)^{3/2}[(a-1)Bz^2(e^{-\lambda_1^*z} + ke^{-\lambda_2^*z}) + az e^{-\lambda_0(z-z_0)}]\}}{d_1 \int_{\mathbb{R}} J_1(y)y^2 e^{\lambda_1^*y} dy};$$

- $p_4 > 0$ large enough such that $z_4 := \{p_4/[(a-1)B]\}^2 > z_0$ and

$$(6.8) \quad p_4 > \frac{8r_2B(a-1) \times \max_{z \geq 0} \{(z+S)^{3/2}[(a-1)Bz^2(e^{-\lambda_2^*z} + he^{-\lambda_1^*z}) + az e^{-\lambda_0(z-z_0)}]\}}{d_2 \int_{\mathbb{R}} J_2(y)y^2 e^{\lambda_2^*y} dy}.$$

It is straightforward to check that the functions $\underline{\phi}_i$ and $\bar{\phi}_i$ are continuous in \mathbb{R} , nonnegative, bounded, of class C^1 except at finitely many points, and that $\underline{\phi}_i \leq \bar{\phi}_i$ in \mathbb{R} , for $i = 1, 2, 3$.

Lemma 6.1. *Under the above assumptions and notations, there exists a positive solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) such that $\underline{\phi}_i \leq \phi_i \leq \bar{\phi}_i$ in \mathbb{R} , $i = 1, 2, 3$.*

Proof. We choose $\varepsilon > 0$ small enough such that

$$(6.9) \quad e^{\lambda_0 z_0} g_3(\lambda_0) + \varepsilon r_3 < 0.$$

Also, up to a translation, condition (2.10) holds for this ε .

We now verify (2.1)-(2.6) for $(\bar{\phi}_1, \bar{\phi}_2, \bar{\phi}_3)$ and $(\underline{\phi}_1, \underline{\phi}_2, \underline{\phi}_3)$ defined in (6.4), for all $z \in \mathbb{R}$ except finitely many points. First of all, it is trivial that (2.1) and (2.2) hold for $z < z_1$ and $z < z_2$ respectively, since $\bar{\phi}_1 = a-1$ in $(-\infty, z_1)$, $\bar{\phi}_2 = a-1$ in $(-\infty, z_2)$, and since $\bar{\phi}_1 \leq a-1$, $\bar{\phi}_2 \leq a-1$, $\bar{\phi}_3 = 1$, $\underline{\phi}_1 \geq 0$ and $\underline{\phi}_2 \geq 0$ in \mathbb{R} . For $z > z_1$, we have $\bar{\phi}_1(z) = (a-1)Bze^{-\lambda_1^* z}$ and, since $z-y > z_1 - S > z_1'$ for all $y \in [-S, S]$, we also have

$$\bar{\phi}_1(z-y) \leq (a-1)B(z-y)e^{-\lambda_1^*(z-y)} \quad \text{for all } y \in [-S, S]$$

(indeed, if $z-y > z_1$, we have equality in the above inequality, and if $z_1' < z-y \leq z_1$, we have $e^{\lambda_1^*(z-y)}/(z-y) \leq B$ and $\bar{\phi}_1(z-y) = a-1 \leq (a-1)B(z-y)e^{-\lambda_1^*(z-y)}$). Hence, for all $z > z_1$, using $J_1 = 0$ almost everywhere outside $[-S, S]$,

$$(6.10) \quad \begin{aligned} \mathcal{N}_1[\bar{\phi}_1](z) &\leq (a-1)B \left[\int_{-S}^S J_1(y)(z-y)e^{-\lambda_1^*(z-y)} dy - ze^{-\lambda_1^* z} \right] \\ &= (a-1)B \left[\int_{\mathbb{R}} J_1(y)(z-y)e^{-\lambda_1^*(z-y)} dy - ze^{-\lambda_1^* z} \right] \\ &= (a-1)Bze^{-\lambda_1^* z} \left[\int_{\mathbb{R}} J_1(y)e^{\lambda_1^* y} dy - 1 \right] - (a-1)Be^{-\lambda_1^* z} \int_{\mathbb{R}} J_1(y)ye^{\lambda_1^* y} dy. \end{aligned}$$

This implies, using $\bar{\phi}_1(z) \geq 0$, $\underline{\phi}_2(z) \geq 0$ and $\bar{\phi}_3(z) = 1$,

$$(6.11) \quad \begin{aligned} \mathcal{U}_1(z) &\leq (a-1)Bze^{-\lambda_1^* z} \{d_1 [I_1(\lambda_1^*) - 1] - s\lambda_1^* + r_1(a-1)\} \\ &\quad - (a-1)Be^{-\lambda_1^* z} \left[d_1 \int_{\mathbb{R}} J_1(y)ye^{\lambda_1^* y} dy - s \right] = 0, \end{aligned}$$

because of (6.3) and $g_1(\lambda_1^*) = 0$. Hence (2.1) holds for all $z \neq z_1$. Similarly, we have $\mathcal{U}_2(z) \leq 0$ for all $z > z_2$, because $g_2(\lambda_2^*) = 0$ and (6.2)-(6.3) hold. Hence (2.2) holds for all $z \neq z_2$. That (2.3) holds for all $z \in \mathbb{R}$ is trivial, since $\alpha(z) \leq 1$, $\bar{\phi}_3(z) = 1$, $\underline{\phi}_1(z) \geq 0$ and $\underline{\phi}_2(z) \geq 0$ for all $z \in \mathbb{R}$.

It remains to check (2.4)-(2.6). Since $\underline{\phi}_1 = 0$ in $(-\infty, z_3]$ and $\underline{\phi}_1 \geq 0$ in \mathbb{R} , (2.4) clearly holds for $z < z_3$. For $z > z_3$ (> 0), we have $\underline{\phi}_1(z) = (a-1)Bze^{-\lambda_1^* z} - p_3 z^{1/2} e^{-\lambda_1^* z}$. Note also from the definitions of $\underline{\phi}_1$ and z_3 that

$$\underline{\phi}_1(x) \geq (a-1)Bxe^{-\lambda_1^* x} - p_3 x^{1/2} e^{-\lambda_1^* x} \quad \text{for all } x > 0.$$

Then, using $z > z_3 > z_0 > z_1 > S$, we have

$$\underline{\phi}_1(z-y)J_1(y) \geq [(a-1)B(z-y)e^{-\lambda_1^*(z-y)} - p_3(z-y)^{1/2}e^{-\lambda_1^*(z-y)}]J_1(y) \quad \text{for all } y \in [-S, S].$$

Using $J_1(y) = 0$ for almost every $y \notin [-S, S]$, it follows that, for all $z > z_3$,

$$(6.12) \quad \begin{aligned} d_1 \mathcal{N}_1[\underline{\phi}_1](z) + s \underline{\phi}'_1(z) &\geq (a-1)Bze^{-\lambda_1^* z} \{d_1[I_1(\lambda_1^*) - 1] - s\lambda_1^*\} \\ &\quad - (a-1)Be^{-\lambda_1^* z} \left\{ d_1 \int_{\mathbb{R}} J_1(y) y e^{\lambda_1^* y} dy - s \right\} + p_3 d_1 z^{1/2} e^{-\lambda_1^* z} \\ &\quad - p_3 e^{-\lambda_1^* z} \left\{ d_1 \int_{\mathbb{R}} J_1(y) (z-y)^{1/2} e^{\lambda_1^* y} dy - s\lambda_1^* z^{1/2} + \frac{s}{2z^{1/2}} \right\}. \end{aligned}$$

On the other hand, we have, for all $z > z_3$ ($> z_0 > \max\{z_1, z_2\}$),

$$\begin{aligned} &r_1 \underline{\phi}_1(z) [-1 - \underline{\phi}_1(z) - k \bar{\phi}_2(z) + a \underline{\phi}_3(z)] \\ &\geq r_1(a-1) \underline{\phi}_1(z) - r_1 \bar{\phi}_1^2(z) - r_1 k \bar{\phi}_1(z) \bar{\phi}_2(z) - r_1 a \bar{\phi}_1(z) e^{-\lambda_0(z-z_0)} \\ &= r_1(a-1)^2 B z e^{-\lambda_1^* z} - r_1(a-1) p_3 z^{1/2} e^{-\lambda_1^* z} - r_1(a-1)^2 B^2 z^2 (e^{-2\lambda_1^* z} + k e^{-(\lambda_1^* + \lambda_2^*)z}) \\ &\quad - r_1 a(a-1) B z e^{-\lambda_1^* z - \lambda_0(z-z_0)}. \end{aligned}$$

Hence we deduce from $g_1(\lambda_1^*) = 0$ and (6.3) that, for all $z > z_3$,

$$\mathcal{L}_1(z) \geq e^{-\lambda_1^* z} [p_3 A_1(z) - A_2(z)],$$

where

$$(6.13) \quad \begin{cases} A_1(z) := d_1 \left[z^{1/2} - \int_{\mathbb{R}} J_1(y) (z-y)^{1/2} e^{\lambda_1^* y} dy \right] + s \lambda_1^* z^{1/2} - \frac{s}{2z^{1/2}} - r_1(a-1) z^{1/2}, \\ A_2(z) := r_1(a-1)^2 B^2 z^2 (e^{-\lambda_1^* z} + k e^{-\lambda_2^* z}) + r_1 a(a-1) B z e^{-\lambda_0(z-z_0)}. \end{cases}$$

Now, using $g_1(\lambda_1^*) = 0$ and (6.3) again, we may rewrite A_1 as

$$A_1(z) = d_1 \int_{\mathbb{R}} J_1(y) \left[z^{1/2} - (z-y)^{1/2} - \frac{y}{2z^{1/2}} \right] e^{\lambda_1^* y} dy = d_1 \int_{-S}^S J_1(y) \left[z^{1/2} - (z-y)^{1/2} - \frac{y}{2z^{1/2}} \right] e^{\lambda_1^* y} dy.$$

Since

$$z^{1/2} - (z-y)^{1/2} - \frac{y}{2z^{1/2}} = \frac{y^2}{2z^{1/2}(z^{1/2} + (z-y)^{1/2})^2} \geq \frac{y^2}{8(z+S)^{3/2}} \quad \text{for } y \in [-S, S],$$

we obtain that

$$(6.14) \quad A_1(z) \geq \frac{d_1}{8(z+S)^{3/2}} \int_{\mathbb{R}} J_1(y) y^2 e^{\lambda_1^* y} dy \quad \text{for all } z > z_3.$$

Therefore, $\mathcal{L}_1(z) \geq 0$ for all $z > z_3$, by the choice of p_3 in (6.7). Hence (2.4) holds for all $z \neq z_3$. Similarly, (2.5) holds for all $z \neq z_4$, using especially $g_2(\lambda_2^*) = 0$ together with (6.3) and (6.8).

Finally, (2.6) holds trivially for $z < z_0$. For $z > z_0$, using $z_0 > \max\{z_1, z_2\} > 0$ together with (1.4), (2.10), (6.6) and $\lambda_0 < \rho$, we have

$$\begin{aligned} \alpha(z) - b \bar{\phi}_1(z) - b \bar{\phi}_2(z) - \underline{\phi}_3(z) &\geq 1 - \varepsilon e^{-\rho z} - b(a-1) B z (e^{-\lambda_1^* z} + e^{-\lambda_2^* z}) - 1 + e^{-\lambda_0(z-z_0)} \\ &\geq -\varepsilon e^{-\rho z} \geq -\varepsilon e^{-\lambda_0 z}. \end{aligned}$$

It follows that, for all $z > z_0$,

$$(6.15) \quad \begin{aligned} \mathcal{L}_3(z) &\geq -e^{-\lambda_0(z-z_0)} \{d_3[I_3(\lambda_0) - 1] - s\lambda_0\} - r_3 \varepsilon [1 - e^{-\lambda_0(z-z_0)}] e^{-\lambda_0 z} \\ &\geq -e^{-\lambda_0 z} [e^{\lambda_0 z_0} g_3(\lambda_0) + r_3 \varepsilon] \geq 0, \end{aligned}$$

due to (6.9). Hence (2.6) holds for all $z \neq z_0$.

Since $0 \leq \underline{\phi}_i \leq \bar{\phi}_i$ in \mathbb{R} for $i = 1, 2, 3$, and each function $\underline{\phi}_i$ is nontrivial, the conclusion of Lemma 6.1 follows from Lemma 2.2, as at the end of the proof of Lemma 4.1. \square

Clearly, the solution (ϕ_1, ϕ_2, ϕ_3) given in Lemma 6.1 satisfies $(\phi_1, \phi_2, \phi_3)(+\infty) = (0, 0, 1)$. Together with Proposition 2.3, Theorem 1.4 follows.

6.2. Waves connecting $(0, 0, 0)$ and $E_1 = (0, 0, 1)$ in the case $s_1^* > s_2^*$: proof of Theorem 1.5. In this subsection, in addition to (6.1), we assume that $s_1^* > s_2^*$, with $s_i^* > 0$ defined in (1.6), and we consider the critical speed

$$s = s_1^*.$$

We recall that $\lambda_1^* > 0$ is the unique minimum of Q_1 in $(0, \hat{\lambda}_1) = (0, +\infty)$. Similarly to (6.3), we also have

$$(6.16) \quad s = s_1^* = d_1 \int_{\mathbb{R}} J_1(y) y e^{\lambda_1^* y} dy.$$

Since $s = s_1^* > s_2^*$, with g_2 given by (4.1) in $[0, \hat{\lambda}_2)$, it then follows from (1.2) and (J1)-(J3) (and the comments after (J1)-(J3)) that there exist $0 < \lambda_3 < \lambda_4 < \hat{\lambda}_2$ such that

$$g_2(\lambda_3) = g_2(\lambda_4) = 0,$$

and $g_2(\lambda) < 0$ if and only if $\lambda \in (\lambda_3, \lambda_4)$. With g_3 given by (4.2) in $[0, \hat{\lambda}_3)$, we have $g_3(0) = 0$ and $g_3'(0) = -s < 0$. Hence we can choose a constant $\lambda_0 > 0$ such that

$$(6.17) \quad 0 < \lambda_0 < \min\{\lambda_1^*, \lambda_3, \rho, \hat{\lambda}_3\} \quad \text{and} \quad g_3(\lambda_0) < 0,$$

where $\rho > 0$ is as in ($\alpha 2$). We also fix a real number μ_3 such that

$$\lambda_3 < \mu_3 < \min\{\lambda_4, \lambda_3 + \lambda_0\}.$$

We then define

$$(6.18) \quad \begin{cases} \bar{\phi}_1(z) = \begin{cases} (a-1)Bze^{-\lambda_1^* z}, & z > z_1, \\ a-1, & z \leq z_1, \end{cases} \\ \underline{\phi}_1(z) = \begin{cases} (a-1)Bze^{-\lambda_1^* z} - p_5 z^{1/2} e^{-\lambda_1^* z}, & z > z_5, \\ 0, & z \leq z_5, \end{cases} \\ \bar{\phi}_2(z) = \min\{(a-1)e^{-\lambda_3 z}, a-1\}, & \underline{\phi}_2(z) = \max\{(a-1)e^{-\lambda_3 z} - p_6 e^{-\mu_3 z}, 0\}, \\ \bar{\phi}_3(z) = 1, & \underline{\phi}_3(z) = \max\{1 - e^{-\lambda_0(z-z_0)}, 0\}, \end{cases}$$

where the parameters $B, z_1, z_0, p_5, z_5, p_6$ are chosen in the following order:

- $B > \lambda_1^* e$ large enough such that $z_1 - z_1' > S$, where S is as in (6.1) and $0 < z_1' < 1/\lambda_1^* < z_1$ are defined by

$$B = \frac{e^{\lambda_1^* z_1}}{z_1} = \frac{e^{\lambda_1^* z_1'}}{z_1'};$$

- $z_0 > z_1$ such that

$$(6.19) \quad b(a-1)Bze^{-\lambda_1^*z} + b(a-1)e^{-\lambda_3z} \leq e^{-\lambda_0(z-z_0)} \quad \text{for all } z \geq z_0$$

(the choice of z_0 is possible since $0 < \lambda_0 < \min\{\lambda_3, \lambda_1^*\}$);

- $p_5 > 0$ large enough such that $z_5 := \{p_5/[(a-1)B]\}^2 > z_0$ and

$$(6.20) \quad p_5 > \frac{8r_1(a-1)B \times \max_{z \geq 0} \{(z+S)^{3/2}[(a-1)Bz^2e^{-\lambda_1^*z} + k(a-1)ze^{-\lambda_3z} + aze^{-\lambda_0(z-z_0)}]\}}{d_1 \int_{\mathbb{R}} J_1(y)y^2 e^{\lambda_1^*y} dy};$$

- $p_6 > 0$ large enough such that $z_6 := (\mu_3 - \lambda_3)^{-1} \ln(p_6/(a-1)) > z_0$ and

$$(6.21) \quad p_6 > \max \left\{ a-1, \frac{r_2(a-1)P_6}{-g_2(\mu_3)} \right\},$$

with

$$P_6 := \max_{z \geq 0} \{h(a-1)Bze^{-(\lambda_1^* + \lambda_3 - \mu_3)z} + (a-1)e^{-(2\lambda_3 - \mu_3)z} + ae^{-\lambda_3z - \lambda_0(z-z_0) + \mu_3z}\}$$

(the choice of p_6 is possible since $\mu_3 < \lambda_3 + \lambda_0 < \min\{2\lambda_3, \lambda_1^* + \lambda_3\}$).

It is straightforward to check that the functions $\underline{\phi}_i$ and $\bar{\phi}_i$ are continuous in \mathbb{R} , nonnegative, bounded, of class C^1 except at finitely many points, and that $\underline{\phi}_i \leq \bar{\phi}_i$ in \mathbb{R} , for $i = 1, 2, 3$.

Lemma 6.2. *Under the above assumptions and notations, there exists a positive solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) such that $\underline{\phi}_i \leq \phi_i \leq \bar{\phi}_i$ in \mathbb{R} , $i = 1, 2, 3$.*

Proof. As in the beginning of the proof of Lemma 6.1, we first choose $\varepsilon > 0$ small enough such that (6.9) holds, as well as condition (2.10), up to a translation.

We now verify (2.1)-(2.6) for $(\bar{\phi}_1, \bar{\phi}_2, \bar{\phi}_3)$ and $(\underline{\phi}_1, \underline{\phi}_2, \underline{\phi}_3)$ defined in (6.18), for all $z \in \mathbb{R}$ except finitely many points. Firstly, it is trivial that (2.1) holds for $z < z_1$, since $\bar{\phi}_1 \leq a-1$, $\bar{\phi}_3 = 1$, $\underline{\phi}_1 \geq 0$ and $\underline{\phi}_2 \geq 0$ in \mathbb{R} . Secondly, it follows from (6.16) and $g_1(\lambda_1^*) = 0$ that (2.1) holds for all $z > z_1$, doing as in (6.10)-(6.11) in the proof of Lemma 6.1. Hence (2.1) holds for all $z \neq z_1$. Thirdly, as in (4.10)-(4.12) in the proof of Lemma 4.1, we have $\mathcal{U}_2(z) \leq 0$, i.e. (2.2), for all $z \neq 0$, using $g_2(\lambda_3) = 0$. Fourthly, that (2.3) holds for all $z \in \mathbb{R}$ is trivial, since $\alpha \leq 1$, $\underline{\phi}_1 \geq 0$ and $\underline{\phi}_2 \geq 0$ in \mathbb{R} .

It remains to show (2.4)-(2.6). Since $\underline{\phi}_1(z) = 0$ for all $z \leq z_5$ and $\underline{\phi}_1 \geq 0$ in \mathbb{R} , (2.4) clearly holds for $z < z_5$. For $z > z_5$ ($> z_0 > 0$), we have $\underline{\phi}_1(z) = (a-1)Bze^{-\lambda_1^*z} - p_5z^{1/2}e^{-\lambda_1^*z}$. Note also from the definitions of $\underline{\phi}_1$ and z_5 that

$$\underline{\phi}_1(x) \geq (a-1)Bxe^{-\lambda_1^*x} - p_5x^{1/2}e^{-\lambda_1^*x} \quad \text{for all } x > 0.$$

Then, using $z > z_5 > z_0 > z_1 > S$, we have

$$\underline{\phi}_1(z-y)J_1(y) \geq [(a-1)B(z-y)e^{-\lambda_1^*(z-y)} - p_5(z-y)^{1/2}e^{-\lambda_1^*(z-y)}]J_1(y) \quad \text{for all } y \in [-S, S].$$

Using $J_1(y) = 0$ for almost every $y \notin [-S, S]$, it follows that, for all $z > z_5$,

$$\begin{aligned} d_1 \mathcal{N}_1[\underline{\phi}_1](z) + s \underline{\phi}'_1(z) &\geq (a-1)Bze^{-\lambda_1^* z} \{d_1[I_1(\lambda_1^*) - 1] - s\lambda_1^*\} \\ &\quad - (a-1)Be^{-\lambda_1^* z} \left\{ d_1 \int_{\mathbb{R}} J_1(y) y e^{\lambda_1^* y} dy - s \right\} + p_5 d_1 z^{1/2} e^{-\lambda_1^* z} \\ &\quad - p_5 e^{-\lambda_1^* z} \left\{ d_1 \int_{\mathbb{R}} J_1(y) (z-y)^{1/2} e^{\lambda_1^* y} dy - s\lambda_1^* z^{1/2} + \frac{s}{2z^{1/2}} \right\}. \end{aligned}$$

On the other hand, for all $z > z_5$ ($> z_0 > z_1 > 0$),

$$\begin{aligned} &r_1 \underline{\phi}_1(z) [-1 - \underline{\phi}_1(z) - k \bar{\phi}_2(z) + a \underline{\phi}_3(z)] \\ &\geq r_1(a-1) \underline{\phi}_1(z) - r_1 \bar{\phi}_1^2(z) - r_1 k \bar{\phi}_1(z) \bar{\phi}_2(z) - r_1 a \bar{\phi}_1(z) e^{-\lambda_0(z-z_0)} \\ &= r_1(a-1)^2 B z e^{-\lambda_1^* z} - r_1(a-1) p_5 z^{1/2} e^{-\lambda_1^* z} \\ &\quad - r_1(a-1)^2 B^2 z^2 e^{-2\lambda_1^* z} - r_1 k (a-1)^2 B z e^{-(\lambda_1^* + \lambda_3)z} - r_1 a (a-1) B z e^{-\lambda_1^* z - \lambda_0(z-z_0)}. \end{aligned}$$

Hence we deduce from $g_1(\lambda_1^*) = 0$ and (6.16) that $\mathcal{L}_1(z) \geq e^{-\lambda_1^* z} [p_5 A_3(z) - A_4(z)]$ for all $z > z_5$, where

$$\begin{cases} A_3(z) := d_1 \left[z^{1/2} - \int_{\mathbb{R}} J_1(y) (z-y)^{1/2} e^{\lambda_1^* y} dy \right] + s \lambda_1^* z^{1/2} - \frac{s}{2z^{1/2}} - r_1(a-1) z^{1/2}, \\ A_4(z) := r_1(a-1)^2 B^2 z^2 e^{-\lambda_1^* z} + r_1 k (a-1)^2 B z e^{-\lambda_3 z} + r_1 a (a-1) B z e^{-\lambda_0(z-z_0)}. \end{cases}$$

Since A_3 has the same expression as A_1 in (6.13), we can proceed as in the proof of (6.14), using $g_1(\lambda_1^*) = 0$ and (6.16) again, and we obtain that

$$A_3(z) \geq \frac{d_1}{8(z+S)^{3/2}} \int_{\mathbb{R}} J_1(y) y^2 e^{\lambda_1^* y} dy \quad \text{for } z > z_5.$$

Then, $\mathcal{L}_1(z) \geq 0$ for $z > z_5$, by the choice of p_5 in (6.20). Hence, (2.4) holds for all $z \neq z_5$.

Now, as in (4.13) in the proof of Lemma 4.1, for every

$$z > z_6 = \frac{1}{\mu_3 - \lambda_3} \ln \left(\frac{p_6}{a-1} \right) > z_0 > z_1 > 0,$$

one has

$$d_2 \mathcal{N}_2[\underline{\phi}_2](z) + s \underline{\phi}'_2(z) \geq (a-1) e^{-\lambda_3 z} \{d_2[I_2(\lambda_3) - 1] - s\lambda_3\} - p_6 e^{-\mu_3 z} \{d_2[I_2(\mu_3) - 1] - s\mu_3\},$$

while

$$\begin{aligned} &r_2 \underline{\phi}_2(z) [-1 - h \bar{\phi}_1(z) - \underline{\phi}_2(z) + a \underline{\phi}_3(z)] \\ &\geq r_2(a-1) \left[(a-1) e^{-\lambda_3 z} - p_6 e^{-\mu_3 z} - h(a-1) B z e^{-(\lambda_1^* + \lambda_3)z} - (a-1) e^{-2\lambda_3 z} - a e^{-\lambda_3 z - \lambda_0(z-z_0)} \right]. \end{aligned}$$

Hence we obtain from $g_2(\lambda_3) = 0$ and (6.21) that

$$\mathcal{L}_2(z) \geq e^{-\mu_3 z} \{-p_6 g_2(\mu_3) - r_2(a-1) P_6\} \geq 0 \quad \text{for all } z > z_6,$$

Together with the fact that (2.5) holds trivially for $z < z_6$, we obtain (2.5) for all $z \neq z_6$.

Finally, (2.6) holds trivially for $z < z_0$. For $z > z_0$, using $z_0 > z_1 > 0$ together with (1.4), (2.10), (6.19) and $\lambda_0 < \rho$, we have

$$\begin{aligned} \alpha(z) - b\bar{\phi}_1(z) - b\bar{\phi}_2(z) - \underline{\phi}_3(z) &\geq 1 - \varepsilon e^{-\rho z} - b(a-1)Bze^{-\lambda_1^* z} - b(a-1)e^{-\lambda_3 z} - 1 + e^{-\lambda_0(z-z_0)} \\ &\geq -\varepsilon e^{-\rho z} \geq -\varepsilon e^{-\lambda_0 z}. \end{aligned}$$

With (6.9), it follows as in (6.15) that $\mathcal{L}_3(z) \geq 0$ for all $z > z_0$. Hence (2.6) holds for all $z \neq z_0$.

Since $0 \leq \underline{\phi}_i \leq \bar{\phi}_i$ in \mathbb{R} for $i = 1, 2, 3$, and each function $\underline{\phi}_1$ is nontrivial, the conclusion of Lemma 6.1 follows from Lemma 2.2, as at the end of the proof of Lemma 4.1. \square

Clearly, the solution (ϕ_1, ϕ_2, ϕ_3) given in Lemma 6.2 satisfies $(\phi_1, \phi_2, \phi_3)(+\infty) = (0, 0, 1)$. Together with Proposition 2.3, Theorem 1.5 follows.

6.3. Waves connecting $(0, 0, 0)$ and $E_2 = (u_p, 0, w_p)$: proof of Theorem 1.6. In this subsection, in addition to (6.1), we assume that $d_1 = d_2 = d_3$, $J_1 = J_2 = J_3$ in \mathbb{R} , that (1.11) is satisfied, and that $\rho \geq \lambda_2^{**}$, where $\rho > 0$ is as in $(\alpha 2)$ and $\lambda_2^{**} > 0$ denotes the unique minimum of the function R_2 defined by (1.9) in $(0, \hat{\lambda}_2) = (0, +\infty)$. We here consider the critical speed

$$s = s_2^{**},$$

where $s_2^{**} > 0$ is defined in (1.9) too. With β_2 and G_2 defined in (1.8) and (5.1) with $s = s_2^{**}$, one has $G_2(\lambda_2^{**}) = 0$, and λ_2^{**} is the unique positive root of this equation. Similarly to (6.3), since $R_2'(\lambda_2^{**}) = 0$, we also have

$$(6.22) \quad s = s_2^{**} = d_2 \int_{\mathbb{R}} J_2(y) y e^{\lambda_2^{**} y} dy.$$

Recalling $A = a - 1 - u_p > 0$ as in (5.3) and $w_p + bu_p = 1$, we then define

$$(6.23) \quad \begin{cases} \bar{\phi}_1(z) = \begin{cases} u_p + ABze^{-\lambda_2^{**} z}, & z > z_1, \\ a - 1, & z \leq z_1, \end{cases} & \underline{\phi}_1(z) = \begin{cases} u_p - u_p Bze^{-\lambda_2^{**} z}, & z > z_1, \\ 0, & z \leq z_1, \end{cases} \\ \bar{\phi}_2(z) = \begin{cases} (a-1)Bze^{-\lambda_2^{**} z}, & z > z_1, \\ a-1, & z \leq z_1, \end{cases} \\ \underline{\phi}_2(z) = \begin{cases} (a-1)Bze^{-\lambda_2^{**} z} - q^* z^{1/2} e^{-\lambda_2^{**} z}, & z > z^*, \\ 0, & z \leq z^*, \end{cases} \\ \bar{\phi}_3(z) = \begin{cases} w_p + bu_p Bze^{-\lambda_2^{**} z}, & z > z_1, \\ 1, & z \leq z_1, \end{cases} & \underline{\phi}_3(z) = \begin{cases} w_p - w_p Bze^{-\lambda_2^{**} z}, & z > z_1, \\ 0, & z \leq z_1, \end{cases} \end{cases}$$

where the parameters B, q^*, z^* are chosen as follows:

- $B > \lambda_2^{**} e$ large enough such that $z_1 - z_2 > S$, where S is as in (6.2) and $0 < z_2 < 1/\lambda_2^{**} < z_1$ are such that

$$B = \frac{e^{\lambda_2^{**} z_1}}{z_1} = \frac{e^{\lambda_2^{**} z_2}}{z_2};$$

- $q^* > 0$ large enough such that $z^* := \{q^*/[(a-1)B]\}^2 > z_1$ and

$$(6.24) \quad q^* > \frac{8r_2(a-1)(2a-1)B^2 \times \max_{z \geq 0} \{(z+S)^{3/2} z^2 e^{-\lambda_2^{**} z}\}}{d_2 \int_{\mathbb{R}} J_2(y) y^2 e^{\lambda_2^{**} y} dy}.$$

It is straightforward to check that the functions $\underline{\phi}_i$ and $\bar{\phi}_i$ are continuous in \mathbb{R} , nonnegative, bounded, of class C^1 except at finitely many points, and that $\underline{\phi}_i \leq \bar{\phi}_i$ in \mathbb{R} , for $i = 1, 2, 3$.

Lemma 6.3. *Under the above assumptions and notations, there exists a positive solution (ϕ_1, ϕ_2, ϕ_3) of (1.3) such that $\underline{\phi}_i \leq \phi_i \leq \bar{\phi}_i$ in \mathbb{R} , $i = 1, 2, 3$.*

Proof. We first choose $\varepsilon > 0$ small enough such that

$$(6.25) \quad \varepsilon \leq \frac{r_2 \beta_2 e}{r_3}.$$

Also, up to a translation, condition (2.10) holds for this ε .

Next, we verify (2.1)-(2.6) for the functions defined in (6.23), for all $z \in \mathbb{R}$ except finitely many points. As before, we only need to show (2.1)-(2.6) for $z > z_1$ or $z > z^*$, respectively: indeed (2.1)-(2.6) hold trivially for $z < z_1$ or $z < z^*$ respectively since $0 \leq \underline{\phi}_1 \leq \bar{\phi}_1 \leq a-1$, $0 \leq \underline{\phi}_2 \leq \bar{\phi}_2 \leq a-1$ and $0 \leq \underline{\phi}_3 \leq \bar{\phi}_3 \leq 1$ in \mathbb{R} , together with the precise definitions (6.23).

For $z > z_1$, $\bar{\phi}_1(z) = u_p + ABze^{-\lambda_2^{**} z}$ and

$$(6.26) \quad \bar{\phi}_1(z-y) \leq u_p + AB(z-y)e^{-\lambda_2^{**}(z-y)} \quad \text{for all } y \in [-S, S]$$

(indeed, $z-y > z_1 - S > z_2$ and if $z-y > z_1$, we have equality in the above inequality, whereas if $z_2 < z-y \leq z_1$, we have $e^{\lambda_2^{**}(z-y)}/(z-y) \leq B$ and $\bar{\phi}_1(z-y) = a-1 = u_p + A \leq u_p + AB(z-y)e^{-\lambda_2^{**}(z-y)}$). Hence

$$\mathcal{N}_1[\bar{\phi}_1](z) \leq ABze^{-\lambda_2^{**} z} [I_1(\lambda_2^{**}) - 1] - ABze^{-\lambda_2^{**} z} \int_{\mathbb{R}} J_1(y) ye^{\lambda_2^{**} y} dy.$$

Also, using $\underline{\phi}_2 \geq 0$ together with $-1 - u_p + aw_p = 0$ and $A = a-1 - u_p = abu_p$, we have

$$-1 - \bar{\phi}_1(z) - k\underline{\phi}_2(z) + a\bar{\phi}_3(z) \leq -1 - u_p - ABze^{-\lambda_2^{**} z} + aw_p + abu_p Bze^{-\lambda_2^{**} z} = 0.$$

This implies that, using $G_2(\lambda_2^{**}) = 0$ and (6.22) together with $d_1 = d_2$ and $J_1 = J_2$,

$$\begin{aligned} \mathcal{U}_1(z) &\leq ABze^{-\lambda_2^{**} z} \{d_1 [I_1(\lambda_2^{**}) - 1] - s\lambda_2^{**}\} - ABze^{-\lambda_2^{**} z} \left[d_1 \int_{\mathbb{R}} J_1(y) ye^{\lambda_2^{**} y} dy - s \right] \\ &= -ABr_2 \beta_2 ze^{-\lambda_2^{**} z} < 0 \end{aligned}$$

for all $z > z_1$. Thus, (2.1) holds for all $z \neq z_1$.

Similar calculations lead to

$$\begin{aligned} \mathcal{U}_2(z) &\leq (a-1)Bze^{-\lambda_2^{**} z} \{d_2 [I_2(\lambda_2^{**}) - 1] - s\lambda_2^{**} + r_2 \beta_2\} \\ &\quad - (a-1)Be^{-\lambda_2^{**} z} \left[d_2 \int_{\mathbb{R}} ye^{-\lambda_2^{**} y} J_2(y) dy - s \right] \\ &\quad + [hu_p - (a-1) + abu_p] r_2 (a-1) B^2 z^2 e^{-2\lambda_2^{**} z} < 0 \end{aligned}$$

for all $z > z_1$, using $G_2(\lambda_2^{**}) = 0$, (6.22) and $hu_p - (a-1) + abu_p < (ab+1)u_p - (a-1) = 0$. Therefore, (2.2) holds for all $z \neq z_1$.

As for (2.3), since $\underline{\phi}_2 \geq 0$ and $1 - bu_p - w_p = 0$, we get that

$$\alpha(z) - b\underline{\phi}_1(z) - b\underline{\phi}_2(z) - \bar{\phi}_3(z) \leq 1 - bu_p + bu_p Bze^{-\lambda_2^{**}z} - w_p - bu_p Bze^{-\lambda_2^{**}z} = 0$$

for every $z > z_1$, and, together with $G_2(\lambda_2^{**}) = 0$ and (6.22), it follows that

$$\begin{aligned} \mathcal{U}_3(z) &\leq bu_p Bze^{-\lambda_2^{**}z} \{d_3[I_3(\lambda_2^{**}) - 1] - s\lambda_2^{**}\} - bu_p Be^{-\lambda_2^{**}z} \left[d_3 \int_{\mathbb{R}} J_3(y) ye^{\lambda_2^{**}y} dy - s \right] \\ &= -bu_p Br_2 \beta_2 ze^{-\lambda_2^{**}z} < 0, \end{aligned}$$

since $J_3 = J_2$ and $d_3 = d_2$. As a consequence, (2.3) holds for all $z \neq z_1$.

For \mathcal{L}_1 and (2.4), one has, for every $z > z_1$, $\underline{\phi}_1(z) = u_p - u_p Bze^{-\lambda_2^{**}z}$ and

$$\underline{\phi}_1(z-y) \geq u_p - u_p B(z-y)e^{-\lambda_2^{**}(z-y)} \quad \text{for all } y \in [-S, S]$$

with the same arguments as for the proof of (6.26), hence

$$\begin{aligned} d_1 \mathcal{N}_1[\underline{\phi}_1](z) + s\underline{\phi}_1(z) &\geq -u_p Bze^{-\lambda_2^{**}z} \{d_1[I_1(\lambda_2^{**}) - 1] - s\lambda_2^{**}\} \\ &\quad + u_p Be^{-\lambda_2^{**}z} \left\{ d_1 \int_{\mathbb{R}} J_1(y) ye^{\lambda_2^{**}y} dy - s \right\} \\ &= u_p Br_2 \beta_2 ze^{-\lambda_2^{**}z} \end{aligned}$$

due to $G_2(\lambda_2^{**}) = 0$, (6.22), $d_1 = d_2$, and $J_1 = J_2$. On the other hand, using $u_p - aw_p = -1$, there holds

$$\begin{aligned} -1 - \underline{\phi}_1(z) - k\bar{\phi}_2(z) + a\underline{\phi}_3(z) &= (-1 - u_p + aw_p) + Bze^{-\lambda_2^{**}z} [u_p - k(a-1) - aw_p] \\ &= -[1 + k(a-1)]Bze^{-\lambda_2^{**}z} \end{aligned}$$

for all $z > z_1$, hence

$$\begin{aligned} \mathcal{L}_1(z) &\geq u_p Br_2 \beta_2 ze^{-\lambda_2^{**}z} - r_1(u_p - u_p Bze^{-\lambda_2^{**}z})[1 + k(a-1)]Bze^{-\lambda_2^{**}z} \\ &\geq u_p Bze^{-\lambda_2^{**}z} \{r_2 \beta_2 - r_1[1 + k(a-1)]\} \geq 0, \end{aligned}$$

using $r_2 \beta_2 \geq r_1[1 + k(a-1)]$ by (1.11). Thus, (2.4) holds for all $z \neq z_1$.

To show (2.5) for $z > z^*$ ($> z_1$), we can proceed for instance as in (6.12) in the proof of Lemma 6.1 to reach

$$\begin{aligned} d_2 \mathcal{N}_2[\underline{\phi}_2](z) + s\underline{\phi}'_2(z) &\geq (a-1)Bze^{-\lambda_2^{**}z} \left\{ d_2 \left[\int_{\mathbb{R}} J_2(y) e^{\lambda_2^{**}y} dy - 1 \right] - s\lambda_2^{**} \right\} \\ &\quad - (a-1)Be^{-\lambda_2^{**}z} \left\{ d_2 \int_{\mathbb{R}} J_2(y) ye^{\lambda_2^{**}y} dy - s \right\} + q^* d_2 e^{-\lambda_2^{**}z} z^{1/2} \\ &\quad - q^* e^{-\lambda_2^{**}z} \left\{ d_2 \int_{\mathbb{R}} J_2(y) (z-y)^{1/2} e^{\lambda_2^{**}y} dy - s\lambda_2^{**} z^{1/2} + \frac{s}{2z^{1/2}} \right\}, \end{aligned}$$

while

$$\begin{aligned}
& r_2 \underline{\phi}_2(z) [-1 - h\bar{\phi}_1(z) - \underline{\phi}_2(z) + a\underline{\phi}_3(z)] \\
&= r_2 \underline{\phi}_2(z) [-1 - hu_p - hABze^{-\lambda_2^{**}z} - \underline{\phi}_2(z) + aw_p - aw_p Bze^{-\lambda_2^{**}z}] \\
&\geq r_2 \beta_2 \underline{\phi}_2(z) - r_2 \bar{\phi}_2(z) (hA + aw_p) Bze^{-\lambda_2^{**}z} - r_2 \bar{\phi}_2^2(z) \\
&\geq r_2 \beta_2 \underline{\phi}_2(z) - r_2 (a-1)(2a-1) B^2 z^2 e^{-2\lambda_2^{**}z},
\end{aligned}$$

using $hA + aw_p + (a-1) < A + aw_p + (a-1) = 2a-1$. It follows from $G_2(\lambda_2^{**}) = 0$ and (6.22) that $\mathcal{L}_2(z) \geq e^{-\lambda_2^{**}z} [q^* A_5(z) - A_6(z)]$ for $z > z^*$, where

$$\begin{cases} A_5(z) := d_2 \left[z^{1/2} - \int_{\mathbb{R}} J_2(y) (z-y)^{1/2} e^{\lambda_2^{**}y} dy \right] + s\lambda_2^{**} z^{1/2} - \frac{s}{2z^{1/2}} - r_2 \beta_2 z^{1/2}, \\ A_6(z) := r_2 (a-1)(2a-1) B^2 z^2 e^{-\lambda_2^{**}z}. \end{cases}$$

Then, using $G_2(\lambda_2^{**}) = 0$ together with (6.22) and (6.24), the same arguments as those in the proofs of Lemmas 6.1 and 6.2 give $\mathcal{L}_2(z) \geq 0$ for all $z > z^*$. Therefore, (2.5) holds for all $z \neq z^*$.

Finally, let us consider \mathcal{L}_3 . For $z > z_1$, $\underline{\phi}_3(z) = w_p - w_p Bze^{-\lambda_2^{**}z}$ and

$$\underline{\phi}_3(z-y) \geq w_p - w_p B(z-y)e^{-\lambda_2^{**}(z-y)} \quad \text{for all } y \in [-S, S],$$

as in the proof of (6.26) for instance. Then, using $d_3 = d_2$ and $J_3 = J_2$, together with $G_2(\lambda_2^{**}) = 0$ and (6.22), one infers that

$$\begin{aligned}
d_3 \mathcal{N}_3[\underline{\phi}_3](z) + s\underline{\phi}_3'(z) &\geq -w_p Bze^{-\lambda_2^{**}z} \{d_3 [I_3(\lambda_2^{**}) - 1] - s\lambda_2^{**}\} + w_p Be^{-\lambda_2^{**}z} \left\{ d_3 \int_{\mathbb{R}} J_3(y) ye^{\lambda_2^{**}y} dy - s \right\} \\
&= w_p Br_2 \beta_2 ze^{-\lambda_2^{**}z}
\end{aligned}$$

for all $z > z_1$. On the other hand, using (2.10), $bu_p + w_p = 1$ and $2b(a-1) < 1$, we obtain

$$\begin{aligned}
& \alpha(z) - b\bar{\phi}_1(z) - b\bar{\phi}_2(z) - \underline{\phi}_3(z) \\
&\geq 1 - \varepsilon e^{-\rho z} - bu_p - bABze^{-\lambda_2^{**}z} - b(a-1)Bze^{-\lambda_2^{**}z} - w_p + w_p Bze^{-\lambda_2^{**}z} \geq -\varepsilon e^{-\rho z}
\end{aligned}$$

for all $z > z_1$, whence

$$\mathcal{L}_3(z) \geq w_p Br_2 \beta_2 ze^{-\lambda_2^{**}z} - r_3 \underline{\phi}_3(z) \varepsilon e^{-\rho z} \geq w_p e^{-\lambda_2^{**}z} (Br_2 \beta_2 z_1 - r_3 \varepsilon) \geq 0,$$

using $\rho \geq \lambda_2^{**}$ and $Br_2 \beta_2 z_1 \geq \lambda_2^{**} \varepsilon r_2 \beta_2 / \lambda_2^{**} = r_2 \beta_2 \varepsilon \geq r_3 \varepsilon$ (because $z_1 > 1/\lambda_2^{**}$, $B > \lambda_2^{**} \varepsilon$ and because of (6.25)). As a consequence, (2.6) holds for all $z \neq z_1$.

Since $0 \leq \underline{\phi}_i \leq \bar{\phi}_i$ in \mathbb{R} for $i = 1, 2, 3$, and each function $\underline{\phi}_i$ is nontrivial, the conclusion of Lemma 6.3 follows from Lemma 2.2, as at the end of the proof of Lemma 4.1. \square

Clearly, the solution (ϕ_1, ϕ_2, ϕ_3) given in Lemma 6.3 satisfies $(\phi_1, \phi_2, \phi_3)(+\infty) = (u_p, 0, w_p)$. Together with Proposition 2.3, Theorem 1.6 follows.

REFERENCES

- [1] H. Berestycki, L. Desvillettes, O. Diekmann, *Can climate change lead to gap formation?* Ecological Complexity 20 (2014), 264-270.
- [2] H. Berestycki, O. Diekmann, C.J. Nagelkerke, P.A. Zegeling, *Can a species keep pace with a shifting climate?* Bull. Math. Biol. 71 (2009), 399-429.
- [3] H. Berestycki, J. Fang, *Forced waves of the Fisher-KPP equation in a shifting environment*, J. Diff. Equations 264 (2018), 2157-2183.
- [4] H. Berestycki, L. Rossi, *Reaction-diffusion equations for population dynamics with forced speed, I - the case of the whole space*, Discrete Contin. Dyn. Syst. 21 (2008), 41-67.
- [5] H. Berestycki, L. Rossi, *Reaction-diffusion equations for population dynamics with forced speed, II - cylindrical type domains*, Discrete Contin. Dyn. Syst. 25 (2009), 19-61.
- [6] J. Bouhours, T. Giletti, *Spreading and vanishing for a monostable reaction-diffusion equation with forced speed*, J. Dynam. Diff. Equations, 31 (2019), 247-286.
- [7] Y.-S. Chen, T. Giletti, J.-S. Guo, *Persistence of preys in a diffusive three species predator-prey system with a pair of strong-weak competing preys*, J. Diff. Equations 281 (2021), 341-378.
- [8] Y.-S. Chen, J.-S. Guo, *Traveling wave solutions for a three-species predator-prey model with two aborigine preys*, Japan J. Industrial Appl. Math. 38 (2021), 455-471.
- [9] Y.-Y. Chen, J.-S. Guo, C.-H. Yao, *Traveling wave solutions for a continuous and discrete diffusive predator-prey model*, J. Math. Anal. Appl. 445 (2017), 212-239.
- [10] W. Choi, T. Giletti, J.-S. Guo, *Persistence of species in a predator-prey system with climate change and either nonlocal or local dispersal*, J. Diff. Equations 302 (2021), 807-853.
- [11] W. Choi, J.-S. Guo, *Forced waves of a three species predator-prey system in a shifting environment*, J. Math. Anal. Appl. 514 (2022), Art. 126283.
- [12] J. Coville, *Can a population survive in a shifting environment using non-local dispersion*, arXiv:2012.09441.
- [13] P. De Leenheer, W. Shen, A. Zhang, *Persistence and extinction of nonlocal dispersal evolution equations in moving habitats*, Nonlinear Analysis: Real World Appl. 54 (2020), Art. 103110.
- [14] F.-D. Dong, B. Li, W.-T. Li, *Forced waves in a Lotka-Volterra competition-diffusion model with a shifting habitat*, J. Diff. Equations, 276 (2021), 433-459.
- [15] A. Ducrot, T. Giletti, J.-S. Guo, M. Shimojo, *Asymptotic spreading speeds for a predator-prey system with two predators and one prey*, Nonlinearity 34 (2021), 669-704.
- [16] A. Ducrot, T. Giletti, H. Matano, *Spreading speeds for multidimensional reaction-diffusion systems of the prey-predator type*, Calc. Var. Part. Diff. Equations 58 (2019), Art. 137.
- [17] A. Ducrot, J.-S. Guo, G. Lin, S. Pan, *The spreading speed and the minimal wave speed of a predator-prey system with nonlocal dispersal*, Z. Angew. Math. Phys. 70 (2019), Art. 146.
- [18] J. Fang, R. Peng, X.-Q. Zhao, *Propagation dynamics of a reaction-diffusion equation in a time-periodic shifting environment*, J. Math. Pures Appl. 147 (2021), 1-28.
- [19] J. Fang, Y. Lou, J. Wu, *Can pathogen spread keep pace with its host invasion?*, SIAM J. Appl. Math. 76 (2016), 1633-1657.
- [20] J.-S. Guo, *Traveling wave solutions for some three-species predator-prey systems*, Tamkang J. Math. 52 (2021), 25-36.
- [21] J.-S. Guo, K.-I. Nakamura, T. Ogiwara, C.-C. Wu, *Traveling wave solutions for a predator-prey system with two predators and one prey*, Nonlinear Analysis: Real World Appl. 54 (2020), Art. 103111.
- [22] F. Hamel, *Reaction-diffusion problems in cylinders with no invariance by translation, Part II: Monotone perturbations*, Ann. Inst. H. Poincaré Anal. Non Linéaire 14 (1997), 555-596.
- [23] H. Hu, X. Zou, *Existence of an extinction wave in the Fisher equation with a shifting habitat*, Proc. Amer. Math. Soc. 145 (2017), 4763-4771.
- [24] M. Kot, *Elements of Mathematical Ecology*, Cambridge Univ. Press, 2003.

- [25] M. Kot, M.A. Lewis, P. van den Driessche, *Dispersal data and the spread of invading organisms*, Ecology 77 (1996), 2027-2042.
- [26] B. Li, S. Bewick, J. Shang, W.F. Fagan, *Persistence and spread of a species with a shifting habitat edge*, SIAM J. Appl. Math. 5 (2014), 1397-1417.
- [27] W.-T. Li, J.-B. Wang, X.-Q. Zhao, *Spatial dynamics of a nonlocal dispersal population model in a shifting environment*, J. Nonlinear Sci. 28 (2018), 1189-1219.
- [28] F. Lutscher, E. Pachepsky, M.A. Lewis, *The effect of dispersal patterns on stream populations*, SIAM J. Appl. Math. 65 (2005), 1305-1327.
- [29] H.-H. Vo, *Persistence versus extinction under a climate change in mixed environments*, J. Diff. Equations 259 (2015), 4947-4988.
- [30] J.-B. Wang, C. Wu, *Forced waves and gap formations for a Lotka-Volterra competition model with nonlocal dispersal and shifting habitats*, Nonlinear Analysis: Real World Appl. 58 (2021), Art. 103208.
- [31] J.-B. Wang, X.-Q. Zhao, *Uniqueness and global stability of forced waves in a shifting environment*, Proc. Amer. Math. Soc. 147 (2019), 1467-1481.
- [32] C.-C. Wu, *On the stable tail limit of traveling wave for a predator-prey system with nonlocal dispersal*, Appl. Math. Letters 113 (2021), Art. 106855.
- [33] C. Wu, Y. Wang, X. Zou, *Spatial-temporal dynamics of a Lotka-Volterra competition model with nonlocal dispersal under shifting environment*, J. Diff. Equations 267 (2019), 4890-4921.
- [34] Y. Yang, C. Wu, Z. Li, *Forced waves and their asymptotics in a Lotka-Volterra cooperative model under climate change*, Appl. Math. Comput. 353 (2019), 254-264.
- [35] Y. Yuan, Y. Wang, X. Zou, *Spatial dynamics of a Lotka-Volterra competition model with a shifting habitat*, Discrete Contin. Dyn. Syst. Ser. B 24 (2019), 5633-5671.
- [36] G.-B. Zhang, W.-T. Li, Z.-C. Wang, *Spreading speeds and traveling waves for nonlocal dispersal equations with degenerate monostable nonlinearity*, J. Diff. Equations 252 (2012), 5096-5124.
- [37] G.-B. Zhang, X.-Q. Zhao, *Propagation dynamics of a nonlocal dispersal Fisher-KPP equation in a time-periodic shifting habitat*, J. Diff. Equations 268 (2020), 2852-2885.
- [38] Y. Zhou, M. Kot, *Life on the move: modeling the effects of climate-driven range shifts with integro-difference equations*, in M.A. Lewis, P.K. Maini, S.V. Petrovskii (Eds.), *Dispersal, individual movement and spatial ecology, a mathematical perspective*, Lecture Notes Math., 2013, pp. 263-292.

DEPARTMENT OF MATHEMATICS, TAMKANG UNIVERSITY, TAMSUI, NEW TAIPEI CITY 251301, TAIWAN
E-mail address: jsguo@mail.tku.edu.tw

AIX MARSEILLE UNIV, CNRS, I2M, MARSEILLE, FRANCE
E-mail address: francois.hamel@univ-amu.fr

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL CHUNG HSING UNIVERSITY, TAICHUNG 402, TAIWAN
E-mail address: chin@email.nchu.edu.tw